

1                    Matrix representations of frame and  
2                    lifted-graphic matroids correspond to gain  
3                    functions

4                    Daryl Funk\*            Daniel Slilaty<sup>†</sup>

5                    November 26, 2018

6                    **Abstract**

7                    Let  $M$  be a 3-connected matroid and let  $\mathbb{F}$  be a field. Let  $A$  be a  
8                    matrix over  $\mathbb{F}$  representing  $M$  and let  $(G, \mathcal{B})$  be a biased graph rep-  
9                    resenting  $M$ . We characterize the relationship between  $A$  and  $(G, \mathcal{B})$ ,  
10                   settling four conjectures of Zaslavsky. We show that for each matrix  
11                   representation  $A$  and each biased graph representation  $(G, \mathcal{B})$  of  $M$ ,  
12                    $A$  is projectively equivalent to a canonical matrix representation arising  
13                   from  $G$  as a gain graph over  $\mathbb{F}^+$  or  $\mathbb{F}^\times$ . Further, we show that  
14                   the projective equivalence classes of matrix representations of  $M$  are  
15                   in one-to-one correspondence with the switching equivalence classes of  
16                   gain graphs arising from  $(G, \mathcal{B})$ .

17                   **1 Introduction**

18                   Let  $M$  be a matroid. Suppose  $M$  has a representation as a matrix  $A$  over a  
19                   field  $\mathbb{F}$  in the standard way: the columns of  $A$  are indexed by  $E(M)$  such that a  
20                   subset of elements of  $M$  is independent precisely when its columns are linearly  
21                   independent; we write  $M = M(A)$ . Suppose  $M$  also has a representation as a  
22                   biased graph: there is a graph  $G$  with  $E(G) = E(M)$  such that when taking  $\mathcal{B}$   
23                   to be the collection of cycles of  $G$  that are circuits of  $M$ , either (i) a subset of

---

\*Department of Mathematics, Douglas College, New Westminster, British Columbia, Canada. Email: [funkd@douglascollege.ca](mailto:funkd@douglascollege.ca)

<sup>†</sup>Department of Mathematics and Statistics, Wright State University, Dayton OH, USA. Email: [daniel.slilaty@wright.edu](mailto:daniel.slilaty@wright.edu). Work partially supported by a grant from the Simons Foundation #246380.

24 elements of  $M$  is independent precisely when it induces a subgraph containing  
 25 at most one cycle, which is not in  $\mathcal{B}$ , or (ii) a subset of elements of  $M$  is  
 26 independent precisely when it induces a subgraph each of whose components  
 27 contains at most one cycle, none of which are in  $\mathcal{B}$ . In the first case,  $(G, \mathcal{B})$   
 28 is a *lifted-graphic representation* of  $M$ ;  $M$  is a *lifted-graphic* matroid and we  
 29 write  $M = L(G, \mathcal{B})$ . In the second case,  $(G, \mathcal{B})$  is a *frame representation* of  
 30  $M$ ;  $M$  is a *frame* matroid and we write  $M = F(G, \mathcal{B})$ .<sup>1</sup> We characterize the  
 31 relationship between such a matrix  $A$  and biased graph  $(G, \mathcal{B})$  when  $M$  is 3-  
 32 connected. While on the surface there is no obvious reason that there should  
 33 be any relationship between these two types of representation, they are in fact  
 34 inherently linked.

35 Frame matroids are a natural generalization of Dowling geometries, and  
 36 are an important class of matroids. Kahn and Kung [9] showed that the only  
 37 matroid varieties containing 3-connected matroids are those given by projec-  
 38 tive geometries over finite fields and Dowling geometries over finite groups.  
 39 Simple frame matroids linearly representable over a finite field are precisely  
 40 the matroids that are members of both of these varieties. Thus it is perhaps  
 41 not surprising (in retrospect!) that these matroids should have a central role  
 42 in matroid structure theory. Geelen, Gerards, and Whittle [8] have proved the  
 43 following far-reaching generalization of Seymour's decomposition theorem for  
 44 regular matroids: If  $\mathcal{M}$  is a proper minor-closed class of  $\text{GF}(q)$ -representable  
 45 matroids, then any member of  $\mathcal{M}$  of sufficiently high connectivity is either a  
 46 bounded-rank perturbation of a frame matroid, the dual of a bounded-rank  
 47 perturbation of a frame matroid, or is representable over a subfield of  $\text{GF}(q)$ .  
 48 Even deeper, Geelen conjectures [6] that a similar result holds for frame ma-  
 49 troids within the class of matroids not containing a  $U_{a,b}$ -minor.

## 50 1.1 Gain graphs and canonical representations

51 The connection between matrix and biased graph representations is provided  
 52 by *gain graphs*.<sup>2</sup> We show that if a matroid  $M$  has a representation as a matrix

---

<sup>1</sup>More generally, we may say  $M$  has a representation as a biased graph  $(G, \mathcal{B})$  if there is a graph  $G$  with  $E(G) = E(M)$  such that when taking  $\mathcal{B}$  to be the collection of cycles of  $G$  that are circuits of  $M$ , there is a partition  $(\mathcal{L}, \mathcal{F})$  of the collection of cycles of  $G$  that are not in  $\mathcal{B}$  such that every cycle in  $\mathcal{L}$  meets every cycle in  $\mathcal{F}$ , and a subset of elements of  $M$  is independent precisely when it induces a subgraph each of whose components contains at most one cycle, none of which are in  $\mathcal{B}$  and at most one of which is in  $\mathcal{L}$ . It is shown in [1] that if  $M$  has both a representation as a matrix over a field and as a biased graph  $(G, \mathcal{B})$  in this way, then  $(G, \mathcal{B})$  is either a lifted-graphic representation or a frame representation of  $M$ .

<sup>2</sup>also known as *group-labelled graphs*.

53 over a field  $\mathbb{F}$  and  $(G, \mathcal{B})$  is a biased graph representing  $M$ , then a direction  
 54 and an element of  $\mathbb{F}$  may be assigned to each edge of  $G$  so that  $G$  becomes a  
 55 gain graph realizing  $\mathcal{B}$ , over the additive or multiplicative group of  $\mathbb{F}$ . Every  
 56 such gain graph yields a *canonical* matrix representation  $A$  for  $M$  over  $\mathbb{F}$ , as  
 57 follows [18].

58 When the gain graph is over the additive group of  $\mathbb{F}$ ,  $A$  is the matrix  
 59 with rows indexed by  $V(G) \cup v_0$ , where  $v_0 \notin V(G)$ , and columns indexed by  
 60  $E(G)$  in which, if  $e$  is assigned element  $a \in \mathbb{F}$ , has distinct endpoints  $u, v$ ,  
 61 and is directed from  $u$  to  $v$ , then entry  $A_{v_0e} = a$ ,  $A_{ue} = 1$ ,  $A_{ve} = -1$ , and  
 62 all remaining entries in column  $e$  are zero; if  $e$  is a loop then  $A_{v_0e} = a$  and  
 63 all remaining entries in column  $e$  are zero. Thus  $A$  consists of the oriented  
 64 incidence matrix of  $G$  with one additional row containing the elements of  $\mathbb{F}$   
 65 that are assigned to each edge  $e \in E(G)$ ; we say  $A$  is a *canonical lift matrix*.  
 66 When the gain graph is over the multiplicative group of  $\mathbb{F}$ ,  $A$  is the matrix  
 67 with rows indexed by  $V(G)$  and columns indexed by  $E(G)$  in which, if  $e$  is  
 68 assigned element  $a \in \mathbb{F}$ , has distinct endpoints  $u, v$ , and is directed from  $u$  to  
 69  $v$ , then  $A_{ue} = 1$ ,  $A_{ve} = -a$ , and the remaining entries in column  $e$  are zero;  
 70 if  $e$  is a loop incident to vertex  $v$  then  $A_{ve} = 1 - a$  and all other entries in  
 71 column  $e$  are zero. We say  $A$  is a *canonical frame matrix*.

72 An assignment of elements of a group  $\Gamma$  to the oriented edges of a graph  
 73 is a  $\Gamma$ -*gain function*. *Switching*, and when  $\Gamma$  is the additive group of a field,  
 74 *scaling*, are two operations that may be applied to a  $\Gamma$ -gain function to obtain  
 75 a new  $\Gamma$ -gain function on the same graph. These operations re-assign elements  
 76 of  $\Gamma$  to  $E(G)$  in such a way that the collection of cycles in  $\mathcal{B}$  determined by  
 77 the new gain function remains unchanged. Thus the biased graph determined  
 78 by the gain graph before and after a switching or scaling operation remains  
 79 the same. Thus when  $\Gamma$  is the additive or multiplicative group of a field, both  
 80 gain graphs provide both a biased graph representation and a canonical matrix  
 81 representation of the same matroid.

82 As will be clear by the citations in Section 2, the notions of frame ma-  
 83 troids, lifted-graphic matroids, their representations by biased graphs, gain  
 84 graphs, their associated canonical matrix representations, and the operations  
 85 of switching and scaling, are all due to Thomas Zaslavsky, presented in his  
 86 seminal series of papers [14, 16, 17, 18].

## 87 1.2 Main results

88 Switching and scaling naturally partition the set of  $\Gamma$ -gain functions on a graph  
 89 into equivalence classes in which two gain functions are equivalent if one may  
 90 be obtained from the other by switching, or when  $\Gamma$  is the additive group

91 of a field, by switching and scaling. When  $(G, \mathcal{B})$  is a biased graph realized  
 92 by a  $\Gamma$ -gain graph, these are the *switching classes* of  $\Gamma$ -realizations of  $(G, \mathcal{B})$ .  
 93 When the group is the additive group  $\mathbb{F}^+$  or multiplicative group  $\mathbb{F}^\times$  of a field  
 94  $\mathbb{F}$ , it is straightforward to show that gain functions belonging to the same  
 95 switching class yield projectively equivalent canonical matrix representations.  
 96 In [18] Zaslavsky conjectured that the converse also holds. Leaving aside some  
 97 technicalities, Conjectures 2.8 and 4.8 of [18] are essentially as follows.

98 **Conjecture 1** (Zaslavsky [18]). *Let  $G$  be a graph and let  $\mathbb{F}$  be a field. The*  
 99 *canonical frame matrices given by two  $\mathbb{F}^\times$ -gain functions  $\varphi$  and  $\psi$  on  $G$  are*  
 100 *projectively equivalent if and only if  $\varphi$  and  $\psi$  are switching equivalent.*

101 **Conjecture 2** (Zaslavsky [18]). *Let  $G$  be a graph and let  $\mathbb{F}$  be a field. The*  
 102 *canonical lift matrices given by two  $\mathbb{F}^+$ -gain functions  $\varphi$  and  $\psi$  on  $G$  are pro-*  
 103 *jectively equivalent if and only if  $\varphi$  and  $\psi$  are switching-and-scaling equivalent.*

104 A *joint* is an unbalanced loop in a biased graph. It is a property of  
 105 the switching operation that the gain on a joint remains unchanged by ev-  
 106 ery switching operation. Thus one of the technicalities to be dealt with in  
 107 Conjectures 1 and 2 is the following. Let  $\varphi_1$  and  $\varphi_2$  be  $\mathbb{F}^\times$ -gain functions on  
 108 a graph  $G$ . Suppose  $\varphi_1(e) = \varphi_2(e)$  for every edge  $e$  that is not a joint, but  
 109 that there is a joint  $e'$  for which  $\varphi_1(e') \neq \varphi_2(e')$  and neither  $\varphi_1(e')$  nor  $\varphi_2(e')$   
 110 is zero. Since a switching operation can alter neither  $\varphi_1(e')$  nor  $\varphi_2(e')$ ,  $\varphi_1$  and  
 111  $\varphi_2$  are not switching equivalent. Yet clearly the canonical matrices defined  
 112 by  $\varphi_1$  and  $\varphi_2$  are projectively equivalent, since each of their columns repre-  
 113 senting the element  $e'$  may be scaled so that the single nonzero entry in each  
 114 is equal to any element of  $\mathbb{F}^\times$ . Our first main result is that for gain graphs  
 115 representing 3-connected matroids, this issue with gains on loops provides the  
 116 only counterexamples to Conjectures 1 and 2.

117 **Theorem 1.** *Let  $M$  be a 3-connected matroid. Let  $(G, \mathcal{B})$  be a biased graph*  
 118 *representing  $M$  and let  $\mathbb{F}$  be a field.*

- 119 1. *The canonical lift matrices given by two  $\mathbb{F}^+$ -gain functions  $\varphi$  and  $\psi$*   
 120 *realizing  $(G, \mathcal{B})$  are projectively equivalent if and only if  $\varphi$  and  $\psi$  are*  
 121 *switching-and-scaling equivalent up to joints.*
- 122 2. *The canonical frame matrices given by two  $\mathbb{F}^\times$ -gain functions  $\varphi$  and  $\psi$*   
 123 *realizing  $(G, \mathcal{B})$  are projectively equivalent if and only if  $\varphi$  and  $\psi$  are*  
 124 *switching equivalent up to joints.*

125 In fact, we can say more. Theorem 1 follows from the stronger statements of  
 126 our Theorems 5.1 and 5.4. Together these also characterize when a canonical

127 lift-matrix representation and a canonical frame-matrix representation may  
 128 be projectively equivalent. Examples that are not 3-connected for which the  
 129 conclusions of Theorem 1 do not hold are not difficult to construct.

130 Let  $M$  be a linearly representable frame or lifted-graphic matroid. Not  
 131 only may  $M$  have projectively inequivalent matrix representations arising as  
 132 canonical matrix representations from the same biased graph, but there may  
 133 also be different biased graphs representing  $M$ . We say a matrix representation  
 134 is *particular* to the biased graph  $(G, \mathcal{B})$  when it is a canonical matrix arising  
 135 from a gain function realizing  $(G, \mathcal{B})$ . Zaslavsky has conjectured the following.

136 **Conjecture 3** ([16]). *Let  $(G, \mathcal{B})$  be a biased graph, where  $G$  is sufficiently con-*  
 137 *nected, and let  $\mathbb{F}$  be a field. If  $F(G, \mathcal{B})$  (resp.  $L(G, \mathcal{B})$ ) is linearly representable*  
 138 *over  $\mathbb{F}$  then  $F(G, \mathcal{B})$  (resp.  $L(G, \mathcal{B})$ ) has a canonical representation particular*  
 139 *to  $(G, \mathcal{B})$ .*

140 Zaslavsky subsequently further conjectured the following.

141 **Conjecture 4** (Zaslavsky, personal communication). *Let  $(G, \mathcal{B})$  be a biased*  
 142 *graph, where  $G$  is sufficiently connected, and let  $\mathbb{F}$  be a field. Every  $\mathbb{F}$ -rep-*  
 143 *resentation of  $F(G, \mathcal{B})$  (respectively  $L(G, \mathcal{B})$ ) is projectively equivalent to a*  
 144 *canonical representation.*

145 Geelen, Gerards, and Whittle prove Conjecture 4 for 3-connected matroids  
 146 in [7], though the result is not explicitly stated (it appears in the proof of their  
 147 Theorem 1.4). Surprisingly, with the exception of a single special case that  
 148 may occur in rank 2, a result even stronger than Conjectures 3 and 4 holds:

149 **Theorem 2.** *Let  $M$  be a 3-connected matroid of rank greater than two, and*  
 150 *let  $\mathbb{F}$  be a field. Let  $A$  be a matrix over  $\mathbb{F}$  representing  $M$  and let  $(G, \mathcal{B})$  be a*  
 151 *biased graph representing  $M$ . Then  $A$  is projectively equivalent to a canonical*  
 152 *representation particular to  $(G, \mathcal{B})$ .*

153 Theorem 2 follows from the stronger statement of Theorem 5.5. Together  
 154 Theorems 5.1, 5.4, and 5.5 imply our next main result. We need just a few  
 155 more definitions before we can state it precisely. A biased graph is *balanced*  
 156 when all of its cycles are balanced, *almost-balanced* when it is not balanced  
 157 but there is a vertex that is contained in every unbalanced cycle of length at  
 158 least two, and *properly unbalanced* otherwise. A properly unbalanced biased  
 159 graph with no pair of vertex-disjoint unbalanced cycles is *tangled*. If  $(G, \mathcal{B})$   
 160 is balanced, then both  $F(G, \mathcal{B})$  and  $L(G, \mathcal{B})$  are equal to the cycle matroid  
 161  $M(G)$  of  $G$ . Since  $M(G)$  has a projectively unique matrix representation  
 162 over every field, and every gain function realizing a balanced biased graph is

163 switching equivalent to the gain function assigning the group identity to every  
 164 element (this follows from Proposition 2.1), Conjectures 1-4 hold in this case.  
 165 Thus we just need consider almost-balanced and properly unbalanced biased  
 166 graphs. The collection of unbalanced cycles in almost-balanced biased graphs  
 167 is highly structured and well-understood (see [5]). For each almost-balanced  
 168 biased graph  $(G, \mathcal{B})$  there is a family of almost-balanced biased graphs  $\mathcal{R}_{(G, \mathcal{B})}$ ,  
 169 each of which represents the frame matroid  $F(G, \mathcal{B})$ . We denote the unique  
 170 biased graph in  $\mathcal{R}_{(G, \mathcal{B})}$  with the least number of loops by  $(\widehat{G}, \widehat{\mathcal{B}})$ ; this is also  
 171 the unique biased graph in the collection for which  $F(\widehat{G}, \widehat{\mathcal{B}}) = L(\widehat{G}, \widehat{\mathcal{B}})$ .

172 Let  $M$  be a 3-connected matroid with rank greater than two. Let  $\mathbb{F}$  be a  
 173 field and let  $(G, \mathcal{B})$  be a biased graph representing  $M$ . Let  $\mathcal{S}_{\mathbb{F}^\times}(G, \mathcal{B})$  denote  
 174 the collection of switching classes of  $\mathbb{F}^\times$ -gain functions realizing  $(G, \mathcal{B})$ , and  
 175 let  $\mathcal{S}_{\mathbb{F}^+}(G, \mathcal{B})$  denote the collection of switching-and-scaling classes of  $\mathbb{F}^+$ -gain  
 176 functions realizing  $(G, \mathcal{B})$ . For each biased graph representing  $M$  and each  
 177 field  $\mathbb{F}$ , define

$$178 \quad \mathcal{S}_{\mathbb{F}}(G, \mathcal{B}) = \begin{cases} \mathcal{S}_{\mathbb{F}^\times}(G, \mathcal{B}) & \text{if } M = F(G, \mathcal{B}) \neq L(G, \mathcal{B}), \\ \mathcal{S}_{\mathbb{F}^+}(G, \mathcal{B}) & \text{if } M = L(G, \mathcal{B}) \neq F(G, \mathcal{B}), \\ \mathcal{S}_{\mathbb{F}^\times}(G, \mathcal{B}) \cup \mathcal{S}_{\mathbb{F}^+}(G, \mathcal{B}) & \text{if } (G, \mathcal{B}) \text{ is tangled,} \\ \mathcal{S}_{\mathbb{F}^+}(\widehat{G}, \widehat{\mathcal{B}}) & \text{if } (G, \mathcal{B}) \text{ is almost-balanced.} \end{cases}$$

179 **Theorem 3.** *Let  $M$  be a 3-connected matroid of rank greater than two. Let*  
 180  *$\mathbb{F}$  be a field and let  $(G, \mathcal{B})$  be a biased graph representing  $M$ . The projec-*  
 181 *tive equivalence classes of matrices over  $\mathbb{F}$  representing  $M$  are in one-to-one*  
 182 *correspondence with the switching classes of gain functions in  $\mathcal{S}_{\mathbb{F}}(G, \mathcal{B})$ .*

183 The proofs of our main results use an inductive argument. For this purpose  
 184 we determine a small collection of biased graphs, at least one of which must  
 185 occur as a biased topological subgraph in every 2-connected biased graph.  
 186 This investigation yields three results on unavoidable minors and topological  
 187 minors of biased graphs that are of independent interest.

188 The graph obtained from a 3-cycle by replacing each edge with a pair of  
 189 parallel edges is denoted  $2C_3$ . The graph obtained from a 4-cycle by replacing  
 190 each edge in a pair of non-adjacent edges with a pair of parallel edges is  
 191 the *tube graph*, denoted  $2C_4''$ . We show that the collection  $\mathcal{G}_0$  consisting of  
 192 the six biased  $2C_3$ 's with no balanced 2-cycle, the three biased tubes with  
 193 no balanced 2-cycle, and the four biased  $K_4$ 's with no balanced triangle is  
 194 the complete set of minor-minimal, 2-connected, properly unbalanced biased  
 195 graphs. Contraction of a loop is never required to obtain one of these minors;  
 196 such a minor is a *link minor*.

197 **Theorem 4.** *Every 2-connected properly unbalanced biased graph contains a*  
 198 *biased graph in  $\mathcal{G}_0$  as a link minor.*

199 We prove an analogue of Theorem 4 for biased topological subgraphs. Let  
 200  $\mathbf{P}$  denote the biased graph consisting of the triangular prism with just its two  
 201 triangles balanced (Figure 5 on page 23). Let  $\mathcal{T}_0$  be the set of biased graphs  
 202 consisting of those in  $\mathcal{G}_0$  together with  $\mathbf{P}$  and the two biased graphs obtained  
 203 from  $\mathbf{P}$  by contracting 1 and 2 edges, respectively, of the matching in  $\mathbf{P}$  linking  
 204 its two triangles.

205 **Theorem 5.** *Every 2-connected properly unbalanced biased graph contains as*  
 206 *a biased subgraph a subdivision of a biased graph in  $\mathcal{T}_0$ .*

207 Theorems 4 and 5 are useful because switching inequivalence of gain func-  
 208 tions can always be found (in the interesting case that there are no joints) on  
 209 a small minor. This is the content of our final main result. Two biased graphs  
 210 representing the 4-point line  $U_{2,4}$  as a frame or lifted-graphic matroid, denoted  
 211  $U_2$  and  $U_3$ , are shown in Figure 7 on page 25.

212 **Theorem 6.** *Let  $(G, \mathcal{B})$  be a 2-connected, loopless, and properly unbalanced*  
 213 *biased graph. Let  $\Gamma$  be an abelian group and let  $\varphi$  and  $\psi$  be  $\Gamma$ -realizations of*  
 214  *$(G, \mathcal{B})$ . Suppose that  $\varphi$  and  $\psi$  are not switching equivalent; in the case  $\Gamma$  is the*  
 215 *additive group of a field, that  $\varphi$  and  $\psi$  are not switching-and-scaling equivalent.*  
 216 *Then either*

- 217 1.  *$(G, \mathcal{B})$  has a link minor  $(H, \mathcal{S}) \in \mathcal{G}_0$  such that the gain functions induced*  
 218 *by  $\varphi$  and  $\psi$  on  $E(H)$  are not switching equivalent (resp. not switching-*  
 219 *and-scaling equivalent), or*
- 220 2.  *$(G, \mathcal{B})$  has  $U_2$  as a minor and  $U_3$  as a link minor such that the gain*  
 221 *functions induced by  $\varphi$  and  $\psi$  are not switching equivalent (resp. not*  
 222 *switching-and-scaling equivalent) on the 2-cycle of  $U_2$  nor on the theta*  
 223 *subgraph of  $U_3$ .*

224 The remainder of the paper is structured as follows. In Section 2 we provide  
 225 necessary preliminary notions. In Section 3 we prove Theorems 4, 5, and 6 on  
 226 unavoidable minors and unavoidable biased topological subgraphs. In Section  
 227 4 we show that Theorems 1 and 2 hold for the set of unavoidable minors, and  
 228 finally in Section 5 we prove Theorems 1, 2, and 3.

## 2 Preliminaries

We assume that the reader is familiar with matroid theory as in Oxley's standard text [10]. In this section we summarize those notions that are central to the results of this paper or are not standard in the literature, and introduce some required notation.

### 2.1 Graphs and biased graphs

Let  $G$  be a graph. We denote the subgraph of  $G$  induced by a subset  $X \subseteq E(G)$  by  $G[X]$ . The set of vertices of  $G[X]$  is denoted  $V(X)$ . A  $k$ -separation of  $G$  is a partition  $(A, B)$  of  $E(G)$  with  $|A| \geq k$ ,  $|B| \geq k$ , and  $|V(A) \cap V(B)| = k$ . A *vertical  $k$ -separation* of  $G$  is a  $k$ -separation  $(A, B)$  of  $G$  with both  $V(A) - V(B)$  and  $V(B) - V(A)$  non-empty. A graph on at least  $k + 2$  vertices is  $k$ -connected if it has no vertical  $l$ -separation for any  $l < k$ . A graph on  $k + 1$  vertices is said to be  $k$ -connected if it has a spanning complete subgraph. Thus a highly connected graph may contain loops or parallel edges. We often need to distinguish between edges with distinct endpoints and loops; thus an edge with distinct endpoints is a *link*.

A *biased graph* is a pair  $(G, \mathcal{B})$  where  $G$  is a graph and  $\mathcal{B}$  is a collection of cycles of  $G$  with the property that no theta subgraph of  $G$  contains exactly two cycles in  $\mathcal{B}$ ; a *theta graph* is the union of three internally disjoint paths linking a pair of vertices. Such a collection  $\mathcal{B}$  is said to satisfy the *theta property*. Cycles in  $\mathcal{B}$  are *balanced*; cycles not in  $\mathcal{B}$  are *unbalanced*. A biased graph is *balanced* if all cycles are balanced, *unbalanced* if contains an unbalanced cycle and *contrabalanced* if no cycle is balanced. Similarly, a subset of edges or a subgraph is *balanced*, *unbalanced*, or *contrabalanced*, according to whether all, not all, or none, respectively, of the cycles it induces or contains are balanced. A vertex  $v$  is a *balancing vertex* if every unbalanced cycle contains  $v$ . A biased graph  $(G, \mathcal{B})$  is  $k$ -connected if  $G$  is  $k$ -connected. For two biased graphs  $(G, \mathcal{B})$  and  $(H, \mathcal{S})$  an *isomorphism*  $\iota: (G, \mathcal{B}) \rightarrow (H, \mathcal{S})$  consists of an underlying graph isomorphism  $\iota: G \rightarrow H$  that takes  $\mathcal{B}$  to  $\mathcal{S}$ . We sometimes write  $\Omega = (G, \mathcal{B})$  and speak of the biased graph  $\Omega$  when there is no need to be explicit about the underlying graph  $G$  and its collection of balanced cycles  $\mathcal{B}$ .

We denote the set of all cycles in  $G$  by  $\mathcal{C}(G)$ . Let  $(G, \mathcal{B})$  be a biased graph and  $e$  an edge in  $G$ . Define  $(G, \mathcal{B}) \setminus e = (G \setminus e, \mathcal{B}|_{G \setminus e})$  where  $\mathcal{B}|_{G \setminus e} = \mathcal{B} \cap \mathcal{C}(G \setminus e)$ . If  $e$  is a link, then define  $(G, \mathcal{B})/e = (G/e, \mathcal{B}|_{G/e})$  where  $\mathcal{B}|_{G/e} = \{C \in \mathcal{C}(G/e) : C \in \mathcal{B} \text{ or } C \cup e \in \mathcal{B}\}$ . If  $e$  is a balanced loop, then  $(G, \mathcal{B})/e = (G, \mathcal{B}) \setminus e$ . In order that contraction of a joint  $e$  of  $(G, \mathcal{B})$  remain consistent with the operation of contraction of  $e$  in the lifted-graphic or frame matroid represented by






Figure 1: Circuits of the frame matroid.

266  $(G, \mathcal{B})$ , two different contraction operations in  $(G, \mathcal{B})$  are required, depending  
 267 upon which matroid  $(G, \mathcal{B})/e$  is to represent. Let  $e$  be a joint of  $(G, \mathcal{B})$ , in-  
 268 cident to vertex  $v$ . To obtain a biased graph representing  $L(G, \mathcal{B})/e$ , define  
 269  $(G, \mathcal{B})/e = (G \setminus e, \mathcal{C}(G \setminus e))$ . To obtain a biased graph representing  $F(G, \mathcal{B})/e$ ,  
 270 define  $(G, \mathcal{B})/e = (G', \mathcal{B}')$  where  $G'$  is obtained from  $G$  by adding each loop  
 271  $e' \neq e$  incident to  $v$  to  $\mathcal{B}$  and replacing each link  $f$  incident to  $v$  with a joint  
 272 incident to its other endpoint. The collection  $\mathcal{B}'$  is  $\mathcal{B}$  restricted to the subgraph  
 273  $G - v$  along with any new balanced loops incident to  $v$ . Whenever contracting  
 274 a joint, we will be explicit about which contraction operation is used, *lift-type*  
 275 or *frame-type*, respectively. In fact, we will only require contraction of a joint  
 276 once; this will be a frame-type contraction. All other minors we consider may  
 277 be obtained without contracting joints. Such a minor is a *link minor*. We  
 278 permit deletion of isolated vertices and do so without comment.

## 279 2.2 Matroids arising from biased graphs

280 A *frame matroid* is a matroid  $M$  to which a basis  $B$  may be added such that  
 281 each  $e \in E(M)$  is contained in the closure of some 2-element subset of  $B$ . A  
 282 subset  $C \subseteq E(G)$  is a circuit of the frame matroid  $F(G, \mathcal{B})$  precisely when  
 283  $C \in \mathcal{B}$  or  $C$  induces a subdivision of one of the graphs shown in Figure 1  
 284 containing no balanced cycle [16]. A matroid  $M$  is *lifted-graphic* if there is a  
 285 single-element extension  $M_0$  of  $M$  by an element  $e_0$  such that  $M_0/e_0$  is graphic.  
 286 Dualizing Crapo's characterization [4] of single-element extensions of matroids,  
 287 Zaslavsky observed [16] that if  $G$  is a graph such that  $M_0/e_0 = M(G)$  then  
 288  $M$  has the natural description in terms of the biased graph  $(G, \mathcal{B})$  of case (i)  
 289 given in the introduction. A subset  $C \subseteq E(G)$  is a circuit of the lifted-graphic  
 290 matroid  $L(G, \mathcal{B})$  precisely when  $C \in \mathcal{B}$  or  $C$  induces a subdivision of one of  
 291 the subgraphs shown in Figure 2 containing no balanced cycle [16]. Minors  
 292 of biased graphs and their matroids agree: for any edge  $e$  of a biased graph  
 293  $(G, \mathcal{B})$ ,  $F(G, \mathcal{B}) \setminus e = F((G, \mathcal{B}) \setminus e)$ ,  $F(G, \mathcal{B})/e = F((G, \mathcal{B})/e)$ ,  $L(G, \mathcal{B}) \setminus e =$   
 294  $L((G, \mathcal{B}) \setminus e)$ , for any link  $e$ ,  $L(G, \mathcal{B})/e = L((G, \mathcal{B})/e)$ , and if  $e$  is a joint then  
 295  $L(G, \mathcal{B})/e = M(G \setminus e)$  [16].

296 Evidently whenever  $(G, \mathcal{B})$  is a biased graph with no two vertex-disjoint




Figure 2: Circuits of the lift matroid.

297 unbalanced cycles,  $F(G, \mathcal{B}) = L(G, \mathcal{B})$ . For notational convenience and to  
 298 avoid potential confusion, in the case that  $(G, \mathcal{B})$  does not have a pair of  
 299 vertex-disjoint unbalanced cycles we denote the matroid on ground set  $E(G)$   
 300 that is equal to both  $F(G, \mathcal{B})$  and  $L(G, \mathcal{B})$  by  $M(G, \mathcal{B})$ .

### 301 2.3 Gain graphs

302 A *gain graph* is obtained from a graph by assigning a direction and an element  
 303 of a group (a *gain*) to each edge of  $G$  [14]. Let  $G$  be a graph and let  $\Gamma$  be a  
 304 group. Orient the edges of  $G$  by arbitrarily assigning a direction to each edge:  
 305 for each edge  $e \in E(G)$  choose one of its ends for its *tail*  $\mathbf{t}(e)$ . The other end  
 306 of  $e$  is its *head*  $\mathbf{h}(e)$ ; if  $e$  is a loop then  $\mathbf{h}(e) = \mathbf{t}(e)$ . Let us specify an oriented  
 307 edge  $e$  by the ordered triple  $(e; u, v)$  where  $u = \mathbf{t}(e)$  and  $v = \mathbf{h}(e)$ . Then  
 308  $e^{-1}$  denotes the ordered triple  $(e; v, u)$ . We think of  $e^{-1}$  as the oriented edge  
 309  $e$  traversed in the reverse direction. Denote the collection of ordered triples  
 310  $\{(e; u, v) : e \in E(G) \text{ has ends } u, v\}$ , consisting of edges of  $G$  together with their  
 311 ends, by  $\vec{E}(G)$ . As long as no confusion may arise, we write  $e$  for both the  
 312 edge  $e \in E(G)$  and for an ordered triple  $(e; u, v) \in \vec{E}(G)$  specifying a direction  
 313 of traversal of  $e$ . Let  $\Gamma$  be a group, and let  $\gamma : \vec{E}(G) \rightarrow \Gamma$  be a function  
 314 satisfying the following condition: for each link  $e$ , if  $\Gamma$  is multiplicative then  
 315  $\gamma(e^{-1}) = \gamma(e)^{-1}$  while if  $\Gamma$  is additive then  $\gamma(e^{-1}) = -\gamma(e)$ . Such a function  
 316 is a  $\Gamma$ -*gain function on  $G$* ; the pair  $(G, \gamma)$  is a  $\Gamma$ -*gain graph*.

317 A gain graph  $(G, \gamma)$  naturally gives rise to a biased graph  $(G, \mathcal{B}_\gamma)$ , where the  
 318 membership of each cycle in the collection of balanced cycles  $\mathcal{B}_\gamma$  is determined  
 319 by the gains assigned to its edges, as follows. Let  $W = (v_1, e_1, v_2, \dots, v_n, e_n,$   
 320  $v_{n+1})$ , where for each  $i$ ,  $v_i, v_{i+1}$  are the ends of edge  $e_i$ , be a walk in  $G$ . Define  
 321  $\gamma(W) = \gamma(e_1; v_1, v_2)\gamma(e_2; v_2, v_3) \cdots \gamma(e_n; v_n, v_{n+1})$  when  $\Gamma$  is multiplicative and  
 322  $\gamma(W) = \gamma(e_1; v_1, v_2) + \gamma(e_2; v_2, v_3) + \cdots + \gamma(e_n; v_n, v_{n+1})$  when  $\Gamma$  is additive.  
 323 For each cycle  $C$  of  $G$  choose a closed Eulerian walk  $W_C$  in  $C$ , and define  $C$   
 324 to be *balanced* with respect to  $\gamma$  if  $\gamma(W_C)$  is the identity element of  $\Gamma$ . Let  $\mathcal{B}_\gamma$   
 325 be the collection of cycles in  $G$  that are balanced with respect to  $\gamma$ . Observe  
 326 that if  $W'_C$  is another closed Eulerian walk in  $C$  then  $\gamma(W_C)$  and  $\gamma(W'_C)$  are  
 327 conjugate. Thus  $\mathcal{B}_\gamma$  is well-defined. If  $(G, \mathcal{B})$  is a biased graph and  $\gamma$  is a

328  $\Gamma$ -gain function such that  $\mathcal{B}_\gamma = \mathcal{B}$ , then we say  $\gamma$  realizes  $(G, \mathcal{B})$  and that  $\gamma$  is  
 329 a  $\Gamma$ -realization of  $(G, \mathcal{B})$ .

330 **Switching and scaling.** Given a  $\Gamma$ -gain function  $\gamma$  on  $G$  and a function  
 331  $\eta: V(G) \rightarrow \Gamma$ , define the gain function  $\gamma^\eta$  by  $\gamma^\eta(e) = \eta(\mathbf{t}(e))^{-1} \cdot \gamma(e) \cdot \eta(\mathbf{h}(e))$  if  
 332  $\Gamma$  is multiplicative and by  $\gamma^\eta(e) = -\eta(\mathbf{t}(e)) + \gamma(e) + \eta(\mathbf{h}(e))$  if  $\Gamma$  is additive. The  
 333 function  $\eta$  is a *switching function*. It is straightforward to check that a cycle  $C$   
 334 is balanced with respect to  $\gamma$  if and only if  $C$  is balanced with respect to  $\gamma^\eta$ , so  
 335  $\mathcal{B}_\gamma = \mathcal{B}_{\gamma^\eta}$ . Observe also that for switching functions  $\eta_1$  and  $\eta_2$ ,  $(\gamma^{\eta_1})^{\eta_2} = \gamma^{\eta_1 \eta_2}$   
 336 or  $\gamma^{\eta_1 + \eta_2}$ . Two  $\Gamma$ -gain functions  $\varphi$  and  $\psi$  are *switching equivalent* if there is a  
 337 switching function  $\eta$  such that  $\varphi^\eta = \psi$ . In the case  $\Gamma$  is the additive group of  
 338 a field  $\mathbb{F}$ , we may choose any nonzero element  $a \in \mathbb{F}$  and obtain a new gain  
 339 function  $a\varphi$  defined by  $a \cdot \varphi(e)$  for each edge  $e$ . Clearly  $\mathcal{B}_{a\varphi} = \mathcal{B}_\varphi$ . We say  $a\varphi$   
 340 is obtained from  $\varphi$  by *scaling*. When  $\Gamma$  is the additive group of a field, we say  
 341 two gain functions  $\varphi$  and  $\psi$  are *switching-and-scaling equivalent* if there is a  
 342 switching function  $\eta$  and scalar  $a \in \mathbb{F}^\times$  such that  $a\varphi^\eta = \psi$ . Evidently, for a  
 343 multiplicative group  $\Gamma$  the relation of being switching equivalent partitions the  
 344 collection of  $\Gamma$ -gain functions on a graph into equivalence classes, its *switching*  
 345 *classes*. Similarly, when  $\Gamma$  is the additive group of a field the relation the being  
 346 switching-and-scaling equivalent partitions the collection of  $\Gamma$ -gain functions  
 347 on a graph into its *switching-and-scaling classes*. Propositions 2.1 and 2.2 are  
 348 immediate.

349 **Proposition 2.1.** *Let  $F$  be a forest of a graph  $G$  and let  $\gamma$  be a  $\Gamma$ -gain function*  
 350 *on  $G$ . There is switching function  $\eta$  such that  $\gamma^\eta(e)$  is the identity for all edges*  
 351  *$e$  in  $F$ .*

352 Let  $F$  be a forest of a graph. A gain function  $\gamma$  is said to be  *$F$ -normalized*  
 353 when  $\gamma(e)$  is the identity for all oriented edges  $e$  in  $F$ .

354 **Proposition 2.2.** *Let  $G$  be a graph, let  $F$  a maximal forest of  $G$ , and let  $\varphi$  and*  
 355  *$\psi$  be two  $F$ -normalized  $\Gamma$ -gain functions on  $G$ . Then  $\varphi$  and  $\psi$  are switching*  
 356 *equivalent if and only if  $\varphi = \psi$ . In the case  $\Gamma$  is the additive group of a field  $\mathbb{F}$ ,*  
 357  *$\varphi$  and  $\psi$  are switching-and-scaling equivalent if and only if  $\varphi = a\psi$  for some*  
 358 *scalar  $a \in \mathbb{F}^\times$ .*

359 **Minors and induced gain functions.** Minors of gain graphs are defined  
 360 so that they are consistent with those of their corresponding biased graphs.  
 361 Let  $\Gamma$  be a group, and let  $(G, \gamma)$  be a  $\Gamma$ -gain graph. Every minor  $H$  of  $G$  has an  
 362 *induced*  $\Gamma$ -gain function  $\gamma|_H$  inherited from  $\gamma$ . Moreover, whenever  $(H, \mathcal{S})$  is a  
 363 biased graph that is a minor of  $(G, \mathcal{B}_\gamma)$  then  $(H, \mathcal{S})$  is realized by the induced

364 gain function  $\gamma|_H$  on  $E(H)$  inherited from  $\gamma$ . We now define these notions and  
 365 justify this claim.

366 Let  $e$  be an edge of  $G$ . Denote by  $(G, \gamma) \setminus e$  and  $(G, \gamma)/e$  the gain graphs  
 367 obtained by deletion and contraction of  $e$  with their induced gain functions  
 368 defined as follows. The induced gain function  $\gamma|_{G \setminus e}$  on  $G \setminus e$  is the restriction  
 369 of  $\gamma$  to  $E(G) - e$ . If  $e$  is a loop assigned the identity element of  $\Gamma$  by  $\gamma$ , then  
 370  $G/e = G \setminus e$  so again the induced gain function is just the restriction of  $\gamma$  to  
 371  $E(G) - e$ . If  $e$  is a link, then there is switching function  $\eta$  such that  $\gamma^\eta(e)$  is  
 372 the identity element of  $\Gamma$ . Define the induced gain function  $\gamma|_{G/e}$  on  $G/e$  to  
 373 be the restriction of  $\gamma^\eta$  to  $E(G) - e$ . Finally, suppose  $e$  is a loop with  $\gamma(e)$  not  
 374 equal to the identify element of  $\Gamma$ . Suppose  $e$  is incident to vertex  $v \in V(G)$ .  
 375 For a lift-type contraction of  $e$ ,  $(G, \gamma)/e$  is the gain graph  $(G \setminus e, \iota)$  where  $\iota$  is  
 376 the gain function assigning the identity element of  $\Gamma$  to every edge; declare  $\iota$   
 377 to be the induced gain function on  $(G, \gamma)/e$ . For a frame-type contraction of  $e$ ,  
 378  $(G, \gamma)/e$  is the gain graph  $(G', \gamma|_{G'})$  in which  $G'$  is the underlying graph of the  
 379 biased graph  $(G, \mathcal{B}_\gamma)/e$  obtained by the frame-type contraction of the joint  $e$   
 380 defined in Section 2.1 above, and  $\gamma|_{G'}$  is the gain function whose restriction to  
 381  $E(G - v)$  is equal to the restriction of  $\gamma$  to  $E(G - v)$ , that assigns the identity  
 382 element of  $\Gamma$  to each balanced loop of  $(G, \mathcal{B}_\gamma)/e$ , and assigns  $\gamma(e)$  to each new  
 383 joint of  $(G, \mathcal{B}_\gamma)/e$ .

384 For link minors, induced gain functions can be defined globally (up to  
 385 switching) as follows. Consider a biased graph  $(G, \mathcal{B})$  and two subsets  $K, D \subseteq$   
 386  $E(G)$  where  $K$  does not contain any loops. Let  $K' \subseteq K$  be a set of edges  
 387 that induce a maximal forest in  $G[K]$ , and let  $D' = D \cup (K \setminus K')$ . Then  
 388  $(G, \mathcal{B})/K \setminus D = (G, \mathcal{B})/K' \setminus D'$ . Thus a link minor may always be obtained  
 389 by contraction of an acyclic set. Consider a gain graph  $(G, \gamma)$  with subsets  
 390  $K, D \subseteq E(G)$  where  $K$  does not contain a loop. We obtain an induced gain  
 391 function  $\gamma|_{G/K \setminus D}$  for  $G/K \setminus D$  as follows. Choose a subset  $K' \subseteq K$  such that  
 392  $G[K']$  is a maximal forest contained in  $G[K]$ , and choose a maximal forest  $F$   
 393 of  $G$  containing  $K'$ . Let  $\gamma^\eta$  be the  $F$ -normalization of  $\gamma$ . Define  $\gamma|_{G/K \setminus D}$  to be  
 394 the restriction of  $\gamma^\eta$  to  $E(G) - (K \cup D)$ .

395 **Proposition 2.3.** *Let  $G$  be a graph and let  $F$  be the edge set of a forest*  
 396 *in  $G$ . Let  $\Gamma$  be an abelian group (resp. the additive group of a field). If  $\varphi$*   
 397 *and  $\psi$  are switching inequivalent (resp. switching-and-scaling inequivalent)  $\Gamma$ -*  
 398 *gain functions on  $G$ , then  $\varphi|_{G/F}$  and  $\psi|_{G/F}$  are switching inequivalent (resp.*  
 399 *switching-and-scaling inequivalent).*

400 *Proof.* Extend  $F$  to a maximal forest  $F_m$  in  $G$  and assume that  $\varphi$  and  $\psi$  are  
 401 normalized on  $F_m$ . Since  $\varphi$  and  $\psi$  are switching inequivalent (resp. switching-  
 402 and-scaling inequivalent), certainly  $\varphi \neq \psi$  (resp.  $\varphi \neq a\psi$  for any scalar  $a$ ), and

403 since both are normalized on  $F_m$ , their restrictions to  $E(G) \setminus F_m$  are not equal  
 404 (resp. neither is obtained by scaling the other). Now in  $G/F$ , the induced  
 405 gain functions  $\varphi|_{G/F}$  and  $\psi|_{G/F}$  are normalized on the maximal forest  $F_m \setminus F$   
 406 of  $G/F$  and  $\varphi|_{G/F} \neq \psi|_{G/F}$ . Thus by Proposition 2.2  $\varphi|_{G/F}$  and  $\psi|_{G/F}$  are  
 407 switching inequivalent (resp. switching-and-scaling inequivalent) on  $G/F$ .  $\square$

408 We say that biased graph  $(G, \mathcal{B})$  has an  $(H, \mathcal{S})$ -minor (respectively,  $(H, \mathcal{S})$ -  
 409 link minor) when there is a minor (resp. link minor)  $(G', \mathcal{B}')$  of  $(G, \mathcal{B})$  that  
 410 is isomorphic to  $(H, \mathcal{S})$ . The following proposition now follows immediately  
 411 from the definitions.

412 **Proposition 2.4.** *If  $\gamma$  is a  $\Gamma$ -realization of  $(G, \mathcal{B})$  and  $(G', \mathcal{B}')$  is a minor of*  
 413  *$(G, \mathcal{B})$ , then the induced gain function  $\gamma|_{G'}$  is a  $\Gamma$ -realization of  $(G', \mathcal{B}')$ .*

414 When  $\gamma$  is a  $\Gamma$ -realization of  $(G, \mathcal{B})$  and  $(G', \mathcal{B}')$  is a minor of  $(G, \mathcal{B})$ , we say  
 415 the  $\Gamma$ -realization  $\gamma|_{G'}$  of  $(G', \mathcal{B}_{G'})$  is the *induced  $\Gamma$ -realization* of  $(G', \mathcal{B}_{G'})$ .

416 **Minors and canonical representations.** Given a matroid  $M$  represented  
 417 over a field by a matrix  $A$ , the operation of removing column  $e$  from  $A$  yields a  
 418 matrix representation of  $M \setminus e$  and the operation of applying row operations so  
 419 that column  $e$  contains a unique nonzero element equal to 1 and then removing  
 420 column  $e$  along with the row in which column  $e$  is nonzero yields a matrix  
 421 representation of  $M/e$ . With just a little more care, we may apply these  
 422 usual operations to canonical matrix representations to obtain a canonical  
 423 matrix representation given by the corresponding induced gain function on  
 424 the gain graph minor. For a canonical matrix  $A$ , and column  $e$  of  $A$ , denote  
 425 by  $A \setminus e$  the matrix obtained by removing column  $e$  from  $A$ . Denote by  $A/e$   
 426 a matrix obtained by applying row operations so that column  $e$  contains a  
 427 unique nonzero element equal to 1, say in row  $i$ , and then removing column  $e$   
 428 and row  $i$  from  $A$ , subject to the following. If  $A$  is a canonical lift matrix and  $e$   
 429 is a link, then row  $i$  is not the “gains row”  $v_0$ . If  $A$  is a canonical frame matrix  
 430 and if  $e$  is a joint, then also scale columns so that for each column  $e' \neq e$  with  
 431 a nonzero entry in row  $i$  and with a second nonzero entry in a row  $j \neq i$ , entry  
 432  $A_{je'}$  is equal to  $1 - \gamma(e)$ .

433 Given an  $\mathbb{F}^\times$ -gain function  $\varphi$  on a graph  $G$ , denote by  $A_F(G, \varphi)$  the canon-  
 434 ical frame matrix defined by  $(G, \varphi)$  as described in Section 1.1. Similarly,  
 435 for an  $\mathbb{F}^+$ -gain function  $\psi$  on  $G$ , denote by  $A_L(G, \psi)$  the canonical lift ma-  
 436 trix defined by  $(G, \psi)$ , as described in Section 1.1. The following lemma is a  
 437 straightforward consequence of the definitions.

438 **Lemma 2.5.** *Let  $G$  be a graph, let  $\mathbb{F}$  be a field, and let  $\varphi: \vec{E}(G) \rightarrow \mathbb{F}^\times$  and*  
 439  *$\psi: \vec{E}(G) \rightarrow \mathbb{F}^+$  be gain functions. Let  $e \in E(G)$ .*

- 440 •  $A_F((G, \varphi) \setminus e) = A_F(G, \varphi) \setminus e$  and  $A_L((G, \varphi) \setminus e) = A_F(G, \varphi) \setminus e$ ,
- 441 •  $A_F((G, \varphi)/e) = A_F(G, \varphi)/e$ , where if  $e$  is a joint the contraction operation is of frame-type,
- 442
- 443 •  $A_L((G, \varphi)/e) = A_F(G, \varphi)/e$ , where if  $e$  is a joint the contraction operation is of lift-type.
- 444

445 Evidently, if  $A$  and  $B$  are projectively equivalent matrices over  $\mathbb{F}$ , then so  
 446 too are  $A \setminus e$  and  $B \setminus e$  projectively equivalent, as are  $A/e$  and  $B/e$ .

## 447 2.4 $\Delta$ - $Y$ and $Y$ - $\Delta$ exchanges

448 The operations of  $\Delta$ - $Y$  and  $Y$ - $\Delta$  exchanges in graphs and in matroids are  
 449 well-understood. Here we generalize these operations from graphs to biased  
 450 graphs, and show that an exchange in a biased graph representation agrees  
 451 with that in the matroid. We use these tools in Sections 4 and 5.

452 Let  $X$  be a triangle in a graph  $G$ . The graph obtained via a  $\Delta$ - $Y$  exchange  
 453 replacing  $X$  with a  $K_{1,3}$ -subgraph is denoted  $\Delta_X G$ . Let  $Y$  be a  $K_{1,3}$ -subgraph  
 454 of  $G$ . The graph obtained via a  $Y$ - $\Delta$  exchange replacing  $Y$  with a triangle is  
 455 denoted  $\nabla_Y G$ . We define the operations of  $\Delta$ - $Y$  and  $Y$ - $\Delta$  exchanges on biased  
 456 graphs as follows. Let  $(G, \mathcal{B})$  be a biased graph and let  $X$  be a balanced  
 457 triangle of  $G$ . Define  $\Delta_X \mathcal{B}$  to be the collection of cycles of  $\Delta_X G$  given by

$$458 \quad \{C \in \mathcal{B} : |C \cap X| = 0 \text{ or } 2\} \cup \{C \Delta X : C \in \mathcal{B} \text{ and } |C \cap X| = 1\}$$

459 where  $\Delta$  denotes symmetric difference.

460 Let  $Y$  be a  $K_{1,3}$ -subgraph of a biased graph  $(G, \mathcal{B})$ . Define  $\nabla_Y \mathcal{B}$  to be  
 461 the collection of cycles of  $\nabla_Y G$  consisting of  $Y$  together with the minimal  
 462 nonempty members of the set

$$463 \quad \{C : C \in \mathcal{B} \text{ and } |C \cap Y| = 0 \text{ or } 2\} \cup \{C \Delta Y : C \in \mathcal{B} \text{ and } |C \cap Y| = 2\}.$$

464 The proofs of the following two propositions are straightforward checks.

465 **Proposition 2.6.** *Let  $X$  be a balanced 3-cycle in a biased graph  $(G, \mathcal{B})$ . Then  
 466  $(\Delta_X G, \Delta_X \mathcal{B})$  is a biased graph.*

467 **Proposition 2.7.** *Let  $Y$  be a  $K_{1,3}$ -subgraph of a biased graph  $(G, \mathcal{B})$ . Then  
 468  $(\nabla_Y G, \nabla_Y \mathcal{B})$  is a biased graph.*

469 We denote the biased graph  $(\Delta_X G, \Delta_X \mathcal{B})$  by  $\Delta_X(G, \mathcal{B})$  and the biased  
 470 graph  $(\nabla_Y G, \nabla_Y \mathcal{B})$  by  $\nabla_Y(G, \mathcal{B})$ . The following is a straightforward conse-  
 471 quence of the definitions. It can be proved by comparing flats.

472 **Proposition 2.8.** *Let  $(G, \mathcal{B})$  be a biased graph. Let  $X$  be a balanced triangle*  
 473 *in  $(G, \mathcal{B})$ , and let  $Y$  be a  $K_{1,3}$ -subgraph of  $G$ .*

474 •  $F(\Delta_X(G, \mathcal{B})) = \Delta_X F(G, \mathcal{B})$  and  $L(\Delta_X(G, \mathcal{B})) = \Delta_X L(G, \mathcal{B})$ .

475 •  $F(\nabla_X(G, \mathcal{B})) = \nabla_X F(G, \mathcal{B})$  and  $L(\nabla_X(G, \mathcal{B})) = \nabla_X L(G, \mathcal{B})$ .

476 The projective equivalence classes of matrix representations of a matroid  
 477 are well-behaved under  $\Delta$ - $Y$  exchanges:

478 **Proposition 2.9** (Whittle [12, Lemma 5.7]). *Let  $M'$  be a matroid obtained*  
 479 *from the matroid  $M$  by a single  $\Delta$ - $Y$  exchange, and let  $\mathbb{F}$  be a field.*

480 1.  $M$  is  $\mathbb{F}$ -representable if and only if  $M'$  is  $\mathbb{F}$ -representable.

481 2. The projective equivalence classes of  $\mathbb{F}$ -representations of  $M$  are in one-  
 482 to-one correspondence with the projective equivalence classes of  $\mathbb{F}$ -repre-  
 483 sentations of  $M'$ .

484 Gain functions realizing a biased graph are similarly well-behaved under  
 485  $\Delta$ - $Y$  exchanges. Proposition 2.10 is an analogue of Proposition 2.9.

486 **Proposition 2.10.** *Let  $X$  be a balanced triangle of a biased graph  $(G, \mathcal{B})$ . Let*  
 487  *$\Gamma$  be the multiplicative (resp. additive) group of a field.*

488 1.  $\varphi$  is a  $\Gamma$ -realization of  $(G, \mathcal{B})$  if and only if  $\varphi$  is a  $\Gamma$ -realization of  
 489  $\Delta_X(G, \mathcal{B})$ .

490 2. The switching (resp. switching-and-scaling) classes of  $\Gamma$ -realizations of  
 491  $(G, \mathcal{B})$  are in one-to-one correspondence with the switching (resp. switch-  
 492 ing-and-scaling) classes of  $\Gamma$ -realizations of  $\Delta_X(G, \mathcal{B})$ .

493 *Proof.* (1) Let  $F$  be a maximal forest of  $\Delta_X G$  containing  $X$ . For each edge  $e \in$   
 494  $X$ ,  $F - e$  is a maximal forest of  $G$  that contains two edges of  $X$ . By Proposition  
 495 2.2, every  $\Gamma$ -realization of  $(G, \mathcal{B})$  is switching equivalent (resp. switching-and-  
 496 scaling equivalent) to a unique (resp. unique up to scaling)  $(F - e)$ -normalized  
 497  $\Gamma$ -realization and every  $\Gamma$ -realization of  $\Delta_X(G, \mathcal{B})$  is switching (resp. switching-  
 498 and-scaling) equivalent to a unique (resp. unique up to scaling)  $F$ -normalized  
 499  $\Gamma$ -realization. Since  $X$  is a balanced triangle of  $(G, \mathcal{B})$ , every  $(F - e)$ -normalized  
 500  $\Gamma$ -realization of  $(G, \mathcal{B})$  also has identity gain value on  $e$ . Thus a  $\Gamma$ -realization  
 501 assigning identity gains to each edge in  $X$  is a  $\Gamma$ -realization of  $(G, \mathcal{B})$  if and  
 502 only if it is a  $\Gamma$ -realization of  $\Delta_X(G, \mathcal{B})$ . Thus we obtain a canonical bijection  
 503 between  $\Gamma$ -gain realizations of  $(G, \mathcal{B})$  and  $\Delta_X(G, \mathcal{B})$ , so (2) holds.  $\square$

504 We now show that a  $\Delta$ - $Y$  exchange applied to a canonical representation  
 505 agrees with the exchange applied to its gain graph. Over any field, a matrix  
 506 representing the matroid  $M(K_4)$  is projectively equivalent to the matrix  $I(K_4)$   
 507 shown below.

$$508 \quad I(K_4) = \begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & 0 & -1 & -1 \\ -1 & -1 & -1 & 0 & 0 & 0 \end{pmatrix}$$

509 The first three columns of this matrix represent a  $K_{1,3}$ -subgraph of  $K_4$  and  
 510 the last three columns a triangle of  $K_4$ . Given a matrix  $A'$  such that  $M(A')$   
 511 contains a triangle  $X$ ,  $A'$  is projectively equivalent to a matrix  $A$  in which the  
 512 columns corresponding to  $X$  are as the last three columns of  $I(K_4)$ , perhaps  
 513 with one row omitted or with zero rows added. Let  $\Delta_X A$  denote the matrix  
 514 obtained from  $A$  by replacing the columns for  $X$  with first three columns of  
 515  $I(K_4)$ , where possibly the fourth row is omitted or additional zero rows are  
 516 added. Then  $M(\Delta_X A) = \Delta_X M(A)$  [2]. Similarly if  $M(A')$  contains a triad  
 517  $Y$ , then  $A'$  is projectively equivalent to a matrix  $A$  in which the columns  
 518 corresponding to  $Y$  are as the first three columns of  $I(K_4)$ , possibly with a  
 519 row omitted or additional zero rows. Let  $\nabla_Y A$  denote the matrix  $A$  obtained  
 520 by replacing the columns of  $Y$  with those of the last three columns of  $I(K_4)$ .  
 521 Then  $M(\nabla_Y A) = \nabla_Y M(A)$  [2]. Thus the next fact follows immediately from  
 522 Proposition 2.10.

523 **Proposition 2.11.** *Let  $\mathbb{F}$  be a field and let  $(G, \varphi)$  be a  $\Gamma$ -gain graph, where  
 524  $\Gamma \in \{\mathbb{F}^\times, \mathbb{F}^+\}$ . Let  $X$  be a balanced triangle of  $(G, \mathcal{B}_\varphi)$  and let  $Y$  be a  $K_{1,3}$ -  
 525 subgraph of  $G$ .*

- 526 •  $\Delta_X A_F(G, \varphi)$  is a canonical frame matrix particular to  $(\Delta_X G, \varphi)$
- 527 •  $\nabla_Y A_F(G, \varphi)$  is a canonical frame matrix particular to  $(\nabla_Y G, \varphi)$
- 528 •  $\Delta_X A_L(G, \varphi)$  is a canonical lift matrix particular to  $(\Delta_X G, \varphi)$
- 529 •  $\nabla_Y A_L(G, \varphi)$  is a canonical lift matrix particular to  $(\nabla_Y G, \varphi)$ .

## 530 2.5 Almost-balanced biased graphs: roll-ups

531 Let  $(G, \mathcal{B})$  be an almost-balanced biased graph with the property that after  
 532 deleting its joints it has a unique balancing vertex  $u$ . Let  $J$  be the set of  
 533 joints of  $(G, \mathcal{B})$  that are not incident to  $u$ , and denote by  $\delta(u)$  the set of



534 links incident to  $u$ . It is not difficult to check that the theta property implies  
 535 that for each pair  $e, e' \in \delta(u)$ , either all cycles containing both  $e$  and  $e'$  are  
 536 balanced or all cycles containing both  $e$  and  $e'$  are unbalanced (for details,  
 537 see [5, Section 1]). Observe that just as for a pair  $e, e' \in \delta(u)$  for which  
 538 every cycle containing both  $e$  and  $e'$  is balanced, for every pair  $f, f' \in J$ ,  
 539 every path linking the endpoint of  $f$  with the endpoint of  $f'$  together with  
 540  $f$  and  $f'$  is a circuit of  $F(G, \mathcal{B})$ . Define  $\Sigma_G(u) = \Sigma(u) = \delta(u) \cup J$ . Then  
 541 for each pair of edges  $e_1, e_2 \in \Sigma_G(u)$ , either every minimal path linking the  
 542 endpoints of  $e_1$  and  $e_2$  in  $G - u$  together with  $\{e_1, e_2\}$  forms a circuit in  
 543  $F(G, \mathcal{B})$ , or all such paths together with  $\{e_1, e_2\}$  are independent in  $F(G, \mathcal{B})$ .  
 544 This defines an equivalence relation on  $\Sigma_G(u)$  [5, Lemma 1.4]. We call these  
 545 equivalence classes the *unbalancing classes* of  $\Sigma_G(u)$ . Now consider the biased  
 546 graph  $(\widehat{G}, \widehat{\mathcal{B}})$  obtained from  $(G, \mathcal{B})$  by replacing each joint  $e \in J$  incident to a  
 547 vertex  $v \neq u$  with a  $uv$ -link. Define  $\widehat{\mathcal{B}}$  to be those cycles having intersection of  
 548 size 0 or 2 with each unbalancing class of  $\Sigma_G(u)$ . It is straightforward to check  
 549 by comparing circuits that  $F(\widehat{G}, \widehat{\mathcal{B}}) = F(G, \mathcal{B})$ . Call  $(\widehat{G}, \widehat{\mathcal{B}})$  the *unrolling of*  
 550  $(G, \mathcal{B})$  to  $u$ . If  $J$  is empty, then set  $(\widehat{G}, \widehat{\mathcal{B}}) = (G, \mathcal{B})$ . Observe that  $u$  is a  
 551 balancing vertex of  $(\widehat{G}, \widehat{\mathcal{B}})$ , and that  $\Sigma_{\widehat{G}}(u) = \Sigma_G(u)$ ; that is, the unbalancing  
 552 classes of  $\Sigma_G(u)$  and of  $\Sigma_{\widehat{G}}(u)$  are the same.

553 For each unbalancing class  $U$  of  $\Sigma_{\widehat{G}}(u)$  there is a biased graph  $(G_U, \mathcal{B}_U)$  for  
 554 which  $F(G_U, \mathcal{B}_U) = F(G, \mathcal{B}) = F(\widehat{G}, \widehat{\mathcal{B}})$ , obtained from  $(\widehat{G}, \widehat{\mathcal{B}})$  by replacing  
 555 each link  $e = uv \in U$  with a joint incident to  $v$  (this fact is straightforward  
 556 to check by comparing circuits; it appears in [5, Proposition 2.2]). Call each  
 557 such biased graph a *roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$  from  $u$* . It is a straightforward check that  
 558 for each unbalancing class  $U \in \Sigma_{\widehat{G}}(u)$ ,  $\Sigma_{G_U}(u) = \Sigma_{\widehat{G}}(u)$ . Note that  $J$  is an  
 559 unbalancing class of  $\Sigma_{\widehat{G}}(u)$ , and that  $(G_J, \mathcal{B}_J) = (G, \mathcal{B})$ . Define  $\mathcal{R}_{(G, \mathcal{B})}$  to be  
 560 the set of biased graphs consisting of  $(\widehat{G}, \widehat{\mathcal{B}})$  together with all of its roll-ups.  
 561 Since each of these biased graphs shares precisely the same set of unbalancing  
 562 classes, we may write simply  $\Sigma(u)$  for this set, when it is clear that we are  
 563 considering a biased graph in the collection  $\mathcal{R}_{(G, \mathcal{B})}$ . Thus  $|\mathcal{R}_{(G, \mathcal{B})}| = |\Sigma(u)| + 1$ .  
 564 Finally, observe that since  $(\widehat{G}, \widehat{\mathcal{B}})$  has no pair of vertex-disjoint unbalanced  
 565 cycles,  $L(\widehat{G}, \widehat{\mathcal{B}}) = F(\widehat{G}, \widehat{\mathcal{B}})$ .

566 There is a special case to consider if  $(G, \mathcal{B})$  is balanced after removing its  
 567 set of joints  $J$ . Let  $x$  be a new isolated vertex added to  $V(G)$ . Then  $J$  may  
 568 be unrolled to any vertex of  $G$ , including  $x$ . In the case  $J$  is unrolled to  $x$ ,  
 569 we obtain a balanced biased graph  $(H, \mathcal{C}(H))$ , so  $F(G, \mathcal{B})$  is equal to the cycle  
 570 matroid  $M(H)$  of  $H$ . The reverse operation may be applied to any graph.  
 571 Given a graph  $H$  and a vertex  $x \in V(H)$ , let  $(G, \mathcal{B})$  be the rollup of the set  
 572 of edges incident to  $x$ ; that is, since the set of edges incident to  $x$  is a single

573 unbalancing class, replace each edge  $xv$  with a joint incident to its endpoint  
 574  $v$ . Then  $M(H) = F(G, \mathcal{B})$ .

575 If a biased graph has two distinct balancing vertices, then it has the very  
 576 restricted form described in Proposition 2.12.

577 **Proposition 2.12** (Zaslavsky [13]). *Let  $(G, \mathcal{B})$  be a 2-connected unbalanced*  
 578 *biased graph with two distinct balancing vertices  $x$  and  $y$ . Then  $G$  is a union*  
 579 *of subgraphs  $G_1 \cup \dots \cup G_m$  where for each pair  $i \neq j$ ,  $G_i \cap G_j = \{x, y\}$ , and a*  
 580 *cycle is in  $\mathcal{B}$  if and only if it is contained in a single subgraph  $G_i$ . If  $m \geq 3$*   
 581 *then  $x$  and  $y$  are the only balancing vertices of  $(G, \mathcal{B})$ .*

582 Observe that if a biased graph  $(G, \mathcal{B})$  of the form described in Proposition  
 583 2.12 has a subgraph  $G_i$  with at least two edges, then  $(E(G_i), E(G) - E(G_i))$   
 584 is a 2-separation of both  $L(G, \mathcal{B})$  and  $F(G, \mathcal{B})$ .

## 585 2.6 Full-rank canonical lift-matrix representations

586 Let  $G$  be a graph and let  $\gamma$  be an  $\mathbb{F}^+$ -gain function on  $G$ , for some field  $\mathbb{F}$ .  
 587 The canonical lift matrix  $A_L(G, \gamma)$  consists of the oriented incidence matrix  
 588 of the subgraph of  $G$  induced by its links together with a row  $v_0$  of gains. It  
 589 is sometimes inconvenient that this matrix is not of full rank. Deleting any  
 590 row other than row  $v_0$  yields a matrix representation of  $L(G, \mathcal{B}_\gamma)$  that is of  
 591 full rank. The oriented incidence matrix of  $G$  is recovered from the resulting  
 592 matrix by appending a row equal to the negation of the sum of all rows but  
 593  $v_0$ . Thus whenever it is convenient we may assume that such a canonical lift  
 594 matrix  $A_L(G, \gamma)$  is of full rank, consisting of a row of gains together with, for  
 595 some vertex  $v \in V(G)$ , the oriented incidence matrix of  $G$  induced by its links  
 596 minus row  $v$ . In the case that  $(G, \mathcal{B}_\gamma)$  has a balancing vertex  $u$  after deleting  
 597 its joints, when it is convenient we assume that this missing vertex is  $u$ .

## 598 3 Unavoidable minors

599 In this section we prove Theorems 4, 5, and 6. We show that there is a small  
 600 collection of biased graphs, at least one of which must appear as a minor  
 601 in every 2-connected biased graph. From this collection we obtain a slightly  
 602 larger collection of biased graphs, and show that every 2-connected properly  
 603 unbalanced biased graph contains a subdivision of at least one of these biased  
 604 graphs. We prove an analogous result for 2-connected almost-balanced biased  
 605 graphs, which we require for the proof of Theorem 5.5. Finally, we show that  
 606 inequivalence of gain functions may always be found on a one of small number  
 607 of unavoidable minors.




Figure 3: The graph  $2C_3$  and its six possible classes of balanced cycles not containing a cycle of length two.

608 **3.1 The minor-minimal, 2-connected, properly unbal-**  
 609 **anced biased graphs**

610 Let  $\mathcal{G}_0$  denote the set of minor-minimal biased graphs that are 2-connected and  
 611 properly unbalanced. We first describe 13 biased graphs in  $\mathcal{G}_0$ , then show that  
 612 these 13 biased graphs form the complete set. Recall that the graph obtained  
 613 from a 3-cycle by replacing each edge with a pair of parallel edges is denoted  
 614  $2C_3$ , and the graph obtained from a 4-cycle by replacing each edge in a pair of  
 615 non-adjacent edges with a pair of parallel edges is the tube, denoted  $2C_4''$ . Six  
 616 of the biased graphs in  $\mathcal{G}_0$  have underlying graph  $2C_3$ , three have underlying  
 617 graph  $2C_4''$ , and four have underlying graph  $K_4$ .

618 The set of cycles of the graph  $K_4$  consists of four triangles and three quadri-  
 619 laterals. We denote by  $D_{t,q} = (K_4, \mathcal{B}_{t,q})$  the biased  $K_4$  with exactly  $t$  balanced  
 620 triangles and  $q$  balanced 4-cycles. There are seven biased  $K_4$ 's:  $D_{0,0}$ ,  $D_{0,1}$ ,  $D_{0,2}$ ,  
 621  $D_{0,3}$ ,  $D_{1,0}$ ,  $D_{2,1}$ , and  $D_{4,2}$  [14]. A biased  $K_4$  is properly unbalanced if and only  
 622 if it does not contain a balanced triangle. Thus the properly unbalanced  $K_4$ 's  
 623 are  $D_{0,0}$ ,  $D_{0,1}$ ,  $D_{0,2}$ , and  $D_{0,3}$ .

624 **Proposition 3.1.** *There are six unlabelled properly unbalanced biased  $2C_3$ 's.*

625 *Proof.* A biased graph  $(2C_3, \mathcal{B})$  is properly unbalanced if and only  $\mathcal{B}$  does not  
 626 contain a 2-cycle. Thus by theta property,  $\mathcal{B}$  is a collection of triangles pairwise  
 627 intersecting in at most one edge. There are eight triangles in  $2C_3$ ; any set of  
 628 five contain a pair that intersect in more than one edge. Hence  $\mathcal{B}$  contains at  
 629 most 4 triangles. The possibilities are shown in Figure 3.  $\square$

630 A biased tube is properly unbalanced if and only it has no balanced 2-cycle.  
 631 There are three such tubes, described in Figure 4.




Figure 4: The graph  $2C_4''$  and its three possible classes of balanced cycles not containing a cycle of length two.

632 We now show that the set of biased graphs consisting of the four biased  
 633  $K_4$ 's with no balanced triangle, the six biased  $2C_3$ 's with no balanced 2-cycle,  
 634 and the three biased tubes with no balanced 2-cycle

$$635 \{D_{0,0}, D_{0,1}, D_{0,2}, D_{0,3}, T_0, T_1, T_2, T'_2, T_3, T_4, B_0, B_1, B_2\}$$

636 forms the complete collection  $\mathcal{G}_0$  of minor-minimal, 2-connected, properly un-  
 637 balanced biased graphs.

638 A *subdivision* of a biased graph  $(G, \mathcal{B})$  is a biased graph  $(H, \mathcal{S})$  in which  $H$   
 639 is a subdivision of  $G$  and a cycle  $C$  of  $H$  is in  $\mathcal{S}$  if and only if its corresponding  
 640 cycle  $C'$  of  $G$  is in  $\mathcal{B}$ . Proposition 3.2 follows immediately from Menger's  
 641 Theorem.

642 **Proposition 3.2.** *Let  $(G, \mathcal{B})$  be a 2-connected biased graph. If  $(G, \mathcal{B})$  contains*  
 643 *a vertex-disjoint pair of unbalanced cycles, neither of which is a loop, then*  
 644  *$(G, \mathcal{B})$  contains a subdivision of  $B_0$ ,  $B_1$ , or  $B_2$ .*

645 Thus to prove Theorem 4 it remains just to show that a properly unbal-  
 646 anced biased graph without two vertex-disjoint unbalanced cycles has a link  
 647 minor from  $\mathcal{G}_0$ . A properly unbalanced biased graph with no two vertex-  
 648 disjoint unbalanced cycles is *tangled*. The structure of tangled signed graphs  
 649 was characterized by Slilaty [11] and the structure of tangled biased graphs  
 650 in general was characterized by Chen and Pivotto [3]. The following theorem  
 651 could be proven as a consequence of Chen and Pivotto's work in [3], but the  
 652 direct proof we present here seems no more difficult.

653 **Theorem 3.3.** *Every tangled biased graph contains as a link minor either a*  
 654 *biased  $2C_3$  with no balanced 2-cycle or a biased  $K_4$  with no balanced triangle.*

655 **Lemma 3.4.** *Let  $\Omega$  be a tangled biased graph. Assume  $\Omega$  contains an unbal-*  
 656 *anced cycle  $C$  and a pair of unbalanced cycles  $C_x$  and  $C_y$  such that  $V(C) \cap$*   
 657  *$V(C_x) = \{x\}$ ,  $V(C) \cap V(C_y) = \{y\}$ , and  $x \neq y$ . Then  $C \cup C_x \cup C_y$  contains a*  
 658 *biased  $2C_3$  with no balanced 2-cycle as a link minor.*

659 *Proof.* Since  $\Omega$  is tangled,  $(C_x \cup C_y) - \{x, y\}$  is connected; furthermore, it is  
 660 vertex-disjoint from  $C$  and so balanced. Let  $K$  be the edge set of  $(C_x \cup C_y) -$   
 661  $\{x, y\}$ . Then  $(C \cup C_x \cup C_y)/K$  is a link minor of  $C \cup C_x \cup C_y$  and is a subdivision  
 662 of a biased  $2C_3$  with no balanced 2-cycle. The result follows.  $\square$

663 *Proof of Theorem 3.3.* Let  $\Omega$  be a link-minor-minimal counterexample. Then  
 664  $|V(\Omega)| > 2$  and  $\Omega$  has no joint, else  $\Omega$  would not be tangled. If  $\Omega$  has more  
 665 than one unbalanced block but no two disjoint unbalanced cycles, then  $\Omega$  must  
 666 have a balancing vertex, a contradiction. Hence  $\Omega$  has only one unbalanced  
 667 block. Evidently our desired minor exists in  $\Omega$  if and only if it exists in the  
 668 unbalanced block of  $\Omega$ . Hence by minimality  $\Omega$  is 2-connected. By minimality  
 669 we may also assume that  $\Omega$  has no balanced 2-cycles.

670 *Claim 1.* The underlying graph of  $\Omega$  is simple.

671 *Proof of Claim:* By way of contradiction assume that  $C$  is an unbalanced 2-  
 672 cycle in  $\Omega$  with vertices  $x$  and  $y$ . Thus  $\Omega - x$  contains an unbalanced cycle  $C_y$   
 673 passing through  $y$  and  $\Omega - y$  contains an unbalanced cycle  $C_x$  passing through  
 674  $x$ . By Lemma 3.4,  $C_x \cup C_y \cup C$  contains a biased  $2C_3$  without a balanced  
 675 2-cycle, a contradiction.  $\clubsuit$

676 If  $\Omega$  has just three vertices, then the underlying graph of  $\Omega$  is a triangle;  
 677 however, now  $\Omega$  has a balancing vertex, a contradiction. Now suppose  $\Omega$  has  
 678 exactly four vertices. For any vertex  $v$ ,  $\Omega - v$  must be an unbalanced triangle  
 679 and so  $\Omega$  is a biased  $K_4$  without a balanced triangle, a contradiction. So for  
 680 the remainder of the proof we can assume that  $\Omega$  has at least five vertices.

681 *Claim 2.* For each vertex  $v$ ,  $\Omega - v$  is unbalanced and has a balancing vertex.

682 *Proof of Claim:* Minimality implies that for any vertex  $v$  in  $\Omega$ ,  $\Omega - v$  is not  
 683 tangled. Since  $\Omega - v$  is unbalanced and has no two disjoint unbalanced cycles,  
 684 it must have a balancing vertex.  $\clubsuit$

685 Given an edge  $e$  with endpoints  $x$  and  $y$ , we denote the vertex in  $\Omega/e$   
 686 resulting from the identification of  $x$  and  $y$  by  $v_e$  or  $v_{xy}$ .

687 *Claim 3.* For each edge  $e$ ,  $\Omega/e$  has  $v_e$  as its unique balancing vertex.

688 *Proof of Claim:* By minimality,  $\Omega/e$  is not tangled and has no two vertex-  
 689 disjoint unbalanced cycles and so must therefore have a balancing vertex. If  
 690 this balancing vertex is  $u \neq v_e$ , then every unbalanced cycle of  $\Omega/e$  passes  
 691 through  $u$  and so every unbalanced cycle of  $\Omega$  passes through  $u$  which implies  
 692 that  $u$  is a balancing vertex of  $\Omega$ , a contradiction.  $\clubsuit$

693 *Claim 4.*  $\Omega$  does not have a vertical 2-separation  $(A, B)$  in which  $B$  is balanced.

694 *Proof of Claim:* Suppose, for a contradiction, that  $(A, B)$  is a vertical 2-  
 695 separation in which  $B$  is balanced. Let  $\{x, y\} = V(A) \cap V(B)$ , and let  $e$  be

696 an edge in  $B$  not incident to at least one of  $x$  and  $y$ . By Claim 3,  $\Omega/e$  has  
 697 balancing vertex  $v_e$ . By our choice of  $e$ ,  $(A, B \setminus e)$  is a 2-separation of  $\Omega/e$ ,  
 698 and  $V(A) \cap V(B \setminus e)$  is either  $\{x, y\}$ ,  $\{x, v_e\}$  or  $\{v_e, y\}$ . In any case, since  $B \setminus e$   
 699 is balanced, every unbalanced cycle of  $\Omega/e$  either does not intersect  $B \setminus e$  or  
 700 intersects  $B \setminus e$  in the edges of a path linking the two vertices of  $V(A) \cap V(B \setminus e)$ .  
 701 But this implies that there is  $v \in \{x, y\}$  such that every unbalanced cycle in  
 702  $\Omega$  contains  $v$ . This makes  $v$  a balancing vertex of  $\Omega$ , a contradiction. ♣

703 *Claim 5.*  $\Omega$  is 3-connected.

704 *Proof of Claim:* Suppose that  $\Omega$  has a vertical 2-separation  $(A, B)$  and let  
 705  $\{x, y\} = V(A) \cap V(B)$ . By Claim 4, neither  $A$  nor  $B$  is balanced. Since  $\Omega$   
 706 does not have a balancing vertex both  $\Omega - x$  and  $\Omega - y$  are unbalanced. Let  
 707  $C_x$  be an unbalanced cycle in  $\Omega - x$  and  $C_y$  be an unbalanced cycle in  $\Omega - y$ .  
 708 Without loss of generality  $E(C_x) \subseteq A$  and either  $E(C_y) \subseteq A$  or  $E(C_y) \subseteq B$ . It  
 709 cannot be that  $E(C_y) \subseteq B$  because then  $C_x$  and  $C_y$  would be vertex-disjoint,  
 710 a contradiction. Hence  $E(C_x) \cup E(C_y) \subseteq A$ ; however, since  $B$  is unbalanced  
 711 any unbalanced cycle  $C'$  in  $\Omega[B]$  intersects both vertices  $x$  and  $y$  and so by  
 712 Lemma 3.4,  $C_x \cup C_y \cup C'$  contains a biased  $2C_3$  having no balanced 2-cycle, a  
 713 contradiction. ♣

714 Now let  $e = xy$  be an edge of  $\Omega$  and let  $E_1, \dots, E_m$  be the unbalancing  
 715 classes of edges incident to balancing vertex  $v_{xy}$  in  $\Omega/e$ . We must have that  
 716  $m \geq 2$ . Now let  $E_{x,i}$  be the edges of  $E_i$  that are incident to  $x$  in  $\Omega$  and  $E_{y,i}$  be  
 717 the edges of  $E_i$  that are incident to  $y$  in  $\Omega$ . Since  $\Omega - y$  is unbalanced, at least  
 718 two of  $E_{x,1}, \dots, E_{x,m}$  are nonempty; similarly, at least two of  $E_{y,1}, \dots, E_{y,m}$   
 719 must be nonempty. Let  $X$  be the set of vertices in  $\Omega - y$  adjacent to  $x$ , and let  
 720  $Y$  be the set of vertices in  $\Omega - x$  that are adjacent to  $y$ . Since the underlying  
 721 graph of  $\Omega$  is simple,  $|X| \geq 2$  and  $|Y| \geq 2$ . Now take  $x_1, x_2 \in X$  such that the  
 722  $xx_1$ - and  $xx_2$ -edges are in different sets  $E_{x,1}, \dots, E_{x,m}$ , and take two similarly  
 723 defined  $y_1, y_2 \in Y$ . Since the underlying graph of  $\Omega$  is simple,  $x_1 \neq x_2$  and  
 724  $y_1 \neq y_2$ .

725 *Claim 6.* Vertices  $x_1, x_2, y_1, y_2$  cannot be chosen so that  $\{x_1, x_2\} \cap \{y_1, y_2\} = \emptyset$ .

726 *Proof of Claim:* Since  $\Omega$  is 3-connected, there is a  $x_1x_2$ -path  $P$  in  $\Omega - \{x, y\}$ .  
 727 For  $i \in \{1, 2\}$  let  $e_i$  denote the  $yy_i$ -edge in  $\Omega$ . Since  $v_{yy_i}$  is a balancing vertex  
 728 in  $\Omega/e_i$  (by Claim 3), the path  $P$  must contain  $y_1$  and  $y_2$  and so there is a  
 729  $y_1y_2$ -path  $P'$  properly contained in  $P$  and  $P'$  that avoids both  $x_1$  and  $x_2$ . The  
 730 unbalanced cycle  $C$  formed by  $y, y_1, P', y_2, y$  avoids  $x_1, x_2$ , and  $x$ ; however, this  
 731 yields a contradiction because contracting the  $xx_1$ -edge in  $\Omega$  leaves a biased  
 732 graph with balancing vertex  $v_{xx_1}$ . ♣

733 *Claim 7.* Vertices  $x_1, x_2, y_1, y_2$  cannot be chosen so that  $|\{x_1, x_2\} \cap \{y_1, y_2\}| = 1$ .




Figure 5:  $\mathsf{P}$  has exactly two triangles, which comprise its set of balanced cycles.

734 *Proof of Claim:* By way of contradiction assume that  $|\{x_1, x_2\} \cap \{y_1, y_2\}| = 1$   
 735 where, without loss of generality,  $x_2 = y_1$ . As in the proof of Claim 6, any  
 736  $x_1x_2$ -path  $P$  in  $\Omega - \{x, y\}$  must contain  $y_2$ . Thus there is a  $y_1y_2$ -path  $P'$   
 737 properly contained in  $P$  and avoiding  $x_1$ , a contradiction. ♣


738 By Claims 6 and 7, every choice of  $x_1, x_2, y_1, y_2$  has (without loss of  
 739 generality)  $x_1 = y_1$  and  $x_2 = y_2$ . This implies that  $X = \{x_1, x_2\} = Y =$   
 740  $\{y_1, y_2\}$ , since otherwise we may choose a third vertex in  $X$  or  $Y$  so that  
 741  $|\{x_1, x_2\} \cap \{y_1, y_2\}| \in \{0, 1\}$ . Let  $e'$  and  $e''$  be respectively the  $xx_1$ - and  $xx_2$ -  
 742 edges in  $\Omega$  and let  $f'$  and  $f''$  be the  $yx_1$ - and  $yx_2$ -edges in  $\Omega$ . It cannot be that  
 743  $e'$  and  $f'$  are in the same unbalancing equivalence class  $E_j \in \{E_1, \dots, E_m\}$   
 744 because then  $\Omega - x_2$  would be balanced. Similarly  $e''$  and  $f''$  are not in the  
 745 same equivalence class. Because  $\Omega$  is 3-connected, there is an  $x_1x_2$ -path  $P$  in  
 746  $\Omega - \{x, y\}$ . The subgraph of  $\Omega$  on edges  $E(P) \cup \{e, e', e'', f', f''\}$  is a subdivision  
 747 of  $K_4$  without a balanced triangle, a contradiction. □

### 748 3.2 Unavoidable topological subgraphs

749 As is often the case with graphs, minors are harder to work with than sub-  
 750 graphs. In this section we prove Theorem 5, an analogue of Theorem 4 for  
 751 biased topological subgraphs. We also prove a result on unavoidable biased  
 752 topological subgraphs for almost-balanced biased graphs.

753 Let  $\mathsf{P}$  denote the biased graph whose underlying graph is the triangular  
 754 prism, with just its two triangles balanced (Figure 5). Let  $L$  denote the match-  
 755 ing of three edges linking the two triangles of  $\mathsf{P}$ . Observe that  $\mathsf{P}/L \cong \mathsf{T}_2$ . Let  
 756  $\mathsf{P}_2$  and  $\mathsf{P}_1$ , respectively, be the biased graphs obtained from  $\mathsf{P}$  by contracting  
 757 1 and 2 edges of  $L$ , respectively (so  $\mathsf{P}_2$  has two edges of  $L$  remaining, and  $\mathsf{P}_1$   
 758 has just one edge of  $L$  remaining from  $\mathsf{P}$ ). Set  $\mathcal{T}_0 = \mathcal{G}_0 \cup \{\mathsf{P}, \mathsf{P}_1, \mathsf{P}_2\}$ . Theo-  
 759 rem 5 guarantees that at least one of the biased graphs in  $\mathcal{T}_0$  is unavoidable  
 760 as a biased topological subgraph in 2-connected properly unbalanced biased  
 761 graphs.

762 *Proof of Theorem 5.* If  $\Omega$  is not tangled, then the result follows from Propo-

Figure 6: The graph  $G'$ .

763 sition 3.2. So assume that  $\Omega$  is tangled. By Theorem 3.3,  $\Omega$  contains a link  
 764 minor  $(G, \mathcal{B})$  that is either a biased  $K_4$  with no balanced triangle or a biased  
 765  $2C_3$  with no balanced 2-cycle. In the first case, since  $K_4$  is 3-regular  $\Omega$  contains  
 766 a subdivision of  $(G, \mathcal{B})$ . In the second case, either  $\Omega$  contains as a subgraph a  
 767 subdivision of  $(G, \mathcal{B})$  or  $\Omega$  contains a link minor  $(G', \mathcal{B}')$  that is 2-connected,  
 768 has minimum degree 3, and contains an edge  $e'$  for which  $(G', \mathcal{B}')/e' = (G, \mathcal{B})$ .  
 769 Since  $\Omega$  is tangled,  $G'$  is as shown in Figure 6. It is straightforward to check  
 770 that if  $(G, \mathcal{B}) \not\cong T_2$ , then  $(G', \mathcal{B}')$  contains a  $K_4$  with no balanced triangle,  
 771 and so our desired subdivision. So assume that  $(G, \mathcal{B}) \cong T_2$ . Then either  
 772  $\mathcal{B}'$  consists of a pair of edge-disjoint triangles both avoiding  $e'$  or  $\mathcal{B}'$  consists  
 773 of a pair of 4-cycles each of which contain  $e'$  but are otherwise edge-disjoint.  
 774 In the latter case, we again have a biased  $K_4$  with no balanced triangle as a  
 775 subgraph, and so are done. In the former case,  $(G', \mathcal{B}') \cong P_1$ . Thus either  $\Omega$   
 776 contains a subdivision of  $P_1$  or  $\Omega$  contains as a link minor  $(G'', \mathcal{B}'')$  which has  
 777 minimum degree 3 and an edge  $e''$  for which  $(G'', \mathcal{B}'')/e'' \cong P_1$ . Either  $\mathcal{B}''$  con-  
 778 sists of a pair of 4-cycles sharing just  $e''$  or  $\mathcal{B}''$  consists of a pair of edge-disjoint  
 779 triangles avoiding  $\{e', e''\}$ . Thus either  $(G'', \mathcal{B}'')$  contains as a subdivision a  
 780 biased  $K_4$  with no balanced triangle, in which case we are done, or  $(G'', \mathcal{B}'')$  is  
 781 isomorphic to  $P_2$ . In the latter case, either  $\Omega$  contains a subdivision of  $P_2$  or  
 782  $\Omega$  contains a link minor  $(G''', \mathcal{B}''')$  with minimum degree 3 and an edge  $e'''$  for  
 783 which  $(G''', \mathcal{B}''')/e''' \cong P_2$ . Thus  $\mathcal{B}'''$  either consists of a pair of disjoint trian-  
 784 gles both avoiding  $\{e', e'', e'''\}$  or a pair of 4-cycles sharing just  $e'''$ . But if  $\mathcal{B}'''$   
 785 consists of a pair of 4-cycles, then  $(G''', \mathcal{B}''')$  contains a pair of vertex-disjoint  
 786 unbalanced triangles, contradicting the fact that  $\Omega$  is tangled. Hence it must  
 787 be the case the  $\mathcal{B}'''$  consists of a pair of disjoint triangles, so  $(G''', \mathcal{B}''') \cong P$ .  $\square$

788 We now show that every almost-balanced biased graph containing a con-  
 789 trabalanced theta contains as a biased topological subgraph one of the biased  
 790 graph in following collection. Denote the graph obtained from  $2C_3$  by deleting  
 791 an edge by  $2C_3 \setminus e$ . The graph  $2C_3 \setminus e$  is obtained from the tube  $2C_4''$  by con-  
 792 tracting one of its non-doubled links, so the cycles of  $2C_3 \setminus e$  are in bijective  
 793 correspondence with the cycles of  $2C_4''$ . Thus there are exactly three biased





Figure 7: Two contrabalanced biased graphs representing  $U_{2,4}$ .

794 graphs  $(2C_3 \setminus e, \mathcal{B})$  without a balanced 2-cycle, each obtained as a single-edge  
 795 contraction of  $B_0$ ,  $B_1$ , or  $B_2$  (Figure 4). We denote these biased graphs  $B'_0$ ,  $B'_1$ ,  
 796 and  $B'_2$ , respectively. Recall that  $D_{1,0} = (K_4, \mathcal{B})$  where  $\mathcal{B}$  consists of exactly  
 797 one balanced triangle;  $D_{1,0}$  has a unique balancing vertex.

798 **Proposition 3.5.** *Let  $(G, \mathcal{B})$  be a 2-connected biased graph that contains a*  
 799 *contrabalanced theta subgraph and a unique balancing vertex after removing*  
 800 *joints. Then  $(G, \mathcal{B})$  contains a subdivision of  $D_{1,0}$ ,  $B'_0$ ,  $B'_1$ , or  $B'_2$ .*

801 *Proof.* Denote by  $nK_2$  the graph consisting of two vertices with  $n$  links between  
 802 them. Since  $(G, \mathcal{B})$  contains a contrabalanced theta subgraph, it contains a  
 803 subdivision of  $(nK_2, \emptyset)$  for some  $n \geq 3$ . Let  $K$  be such a subdivision in  $(G, \mathcal{B})$   
 804 with  $n$  as large as possible. Let  $u$  be the balancing vertex of  $(G, \mathcal{B})$ . One of the  
 805 two degree- $n$  vertices of  $K$  is  $u$ ; let  $v$  be the other degree- $n$  vertex of  $K$ . Then  
 806  $K$  is the union of  $n$  internally disjoint  $u$ - $v$ -paths  $P_1, \dots, P_n$ . By assumption  
 807  $(G, \mathcal{B})$  does not have the structure described in Proposition 2.12. Thus there  
 808 is a path  $P$  in  $G$  internally disjoint from  $K$  with both its ends in  $K$  such that  
 809 either


- 810 • both ends of  $P$  are internal vertices of two distinct paths  $P_i$  and  $P_j$ , or
- 811 •  $P$  has  $u$  as one end, an internal vertex of one of the paths  $P_i$  as its other
- 812 end, and the cycle contained in  $P \cup P_i$  is unbalanced.

813 In the first case,  $P_i \cup P_j \cup P$  is a theta subgraph with its cycle  $P_i \cup P_j$  unbalanced  
 814 and its cycle avoiding  $u$  balanced. By the theta property the cycle in  $P_i \cup P_j \cup P$   
 815 avoiding  $v$  is unbalanced. Thus  $K \cup P$  contains a subdivision of  $D_{1,0}$ . In the  
 816 second case,  $K \cup P$  contains a subdivision of  $B'_0$ ,  $B'_1$ , or  $B'_2$ . □

### 817 3.3 Confining inequivalence to a small minor

818 We can now prove Theorem 6.

819 Two biased graphic representations of  $U_{2,4}$  are  $U_2$  and  $U_3$ , shown in Figure  
 820 7; all cycles in each are unbalanced. Theorem 6 localizes switching inequiva-  
 821 lence of gain functions on a small minor: if not a biased graph in  $\mathcal{G}_0$  then on  
 822 both  $U_2$  and  $U_3$ .

Figure 8: The possibilities for  $\Omega$ .

823 *Proof of Theorem 6.* By Theorem 5,  $(G, \mathcal{B})$  has a biased subgraph  $(G_0, \mathcal{B}_0)$   
824 that is a subdivision of a member of  $\mathcal{G}_0 \cup \{\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}\}$ . Since  $G$  is 2-connected and  
825 loopless, there is a sequence of 2-connected subgraphs  $(G_0, \mathcal{B}_0), \dots, (G_n, \mathcal{B}_n)$   
826 such that  $(G_n, \mathcal{B}_n) = (G, \mathcal{B})$  and  $(G_{i+1}, \mathcal{B}_{i+1}) = (G_i, \mathcal{B}_i) \cup P_i$  for some path  
827  $P_i$  that is internally disjoint from  $G_i$ . Let  $\varphi_i$  and  $\psi_i$  be the  $\Gamma$ -gain functions  
828 induced by  $\varphi$  and  $\psi$  on  $G_i$ . If  $\varphi_0$  and  $\psi_0$  are switching inequivalent (resp.,  
829 switching-and-scaling inequivalent in the case  $\Gamma$  is the additive group of a  
830 field), then the result follows by Proposition 2.3. Otherwise, there is an integer  
831  $t \in \{0, \dots, n-1\}$  such that  $\varphi_i$  and  $\psi_i$  are switching (resp., switching-and-  
832 scaling) equivalent for  $i \leq t$  and  $\varphi_{t+1}$  and  $\psi_{t+1}$  are switching (resp., switching-  
833 and-scaling) inequivalent. Let  $e$  be an edge in path  $P_t$ . Since  $G_{t+1}$  is 2-  
834 connected, there is a spanning tree  $T_{t+1}$  of  $G_{t+1}$  that does not contain  $e$ .  
835 Normalize  $\varphi_{t+1}$  and  $\psi_{t+1}$  on  $T_{t+1}$ . Let  $T_t$  be  $T_{t+1}$  restricted to  $G_t$ . Then  $T_t$  is  
836 a spanning tree of  $G_t$ . Since  $\varphi_{t+1}$  and  $\psi_{t+1}$  are normalized on  $T_{t+1}$ ,  $\varphi_t$  and  $\psi_t$   
837 are normalized on  $T_t$ . Since  $\varphi_t$  and  $\psi_t$  are switching (-and-scaling) equivalent  
838 while  $\varphi_{t+1}$  and  $\psi_{t+1}$  are switching (-and-scaling) inequivalent, by Proposition  
839 2.2  $\varphi_t = \psi_t$  (resp.  $\varphi_t = a\psi_t$  for some scalar  $a$ ) while  $\varphi_{t+1}$  and  $\psi_{t+1}$  are equal  
840 (resp. equal up to scaling) everywhere except at  $e$ . Hence for every cycle  $C$   
841 of  $G_{t+1}$  containing path  $P_t$ ,  $\varphi_{t+1}(C) \neq \psi_{t+1}(C)$ . Since  $\varphi_{t+1}$  and  $\psi_{t+1}$  are  $\Gamma$ -  
842 realizations of  $(G_{t+1}, \mathcal{B}_{t+1})$ , it must be that every such cycle  $C$  is unbalanced.  
843 Extend  $P_t$  in  $G_{t+1}$  to a path  $P$  that is internally disjoint from  $G_0$  but whose  
844 endpoints are both on  $G_0$ . Let  $\varphi'$  and  $\psi'$  be  $\varphi_{t+1}$  and  $\psi_{t+1}$  restricted to the  
845 biased graph  $(G_0, \mathcal{B}_0) \cup P$ . Again,  $\varphi'$  and  $\psi'$  are equal (resp. equal up to  
846 scaling) on every edge of  $(G_0, \mathcal{B}_0) \cup P$  save for the edge  $e$  and every cycle  $C$  in  
847  $(G_0, \mathcal{B}_0) \cup P$  containing  $e$  is therefore unbalanced. Now in  $(G_0, \mathcal{B}_0) \cup P$  there  
848 is a link minor  $\Omega = ((G_0, \mathcal{B}_0) \cup P)/K \setminus D$  for which  $\Omega \setminus e$  is in  $\mathcal{G}_0$  or  $\Omega \setminus e/f$  is in  
849  $\mathcal{G}_0$  for some link  $f$ . The possibilities for  $\Omega$  are as shown in Figure 8.

850 Re-normalize  $\varphi'$  and  $\psi'$  on a spanning tree of  $(G_0, \mathcal{B}_0) \cup (P - e)$  that contains  
 851 the contraction set  $K$  and let  $\varphi|_\Omega$  and  $\psi|_\Omega$  be the induced gain functions on  $\Omega$ .  
 852 Again,  $\varphi|_\Omega$  and  $\psi|_\Omega$  are equal (equal up to scaling) on each edge of  $(G_0, \mathcal{B}_0) \cup$   
 853  $(P - e)$  but differ (differ even after scaling) on edge  $e$ , and every cycle containing  
 854  $e$  is unbalanced. The first outcome of the theorem holds in cases (i), (ii), (iv),  
 855 and (v) of Figure 8 while the second outcome holds in cases (iii) and (vi). A  
 856 single frame-type contraction of a joint is necessary in order to obtain  $\mathbf{U}_2$  in  
 857 case (vi), but no other contraction of a joint is required. Thus all minors but  
 858 this one are link minors.  $\square$

859 The following observation will be used in the proof of Theorem 5.1.

860 **Corollary 3.6.** *If  $(G, \mathcal{B})$  is tangled, then the first outcome of Theorem 6 holds.*

861 *Proof.* If the first outcome of Theorem 6 does not hold, then the biased graph  
 862  $\Omega$  in the proof of Theorem 6 is that of either case (iii) or (vi) of Figure 8. Each  
 863 of these contain a pair of vertex-disjoint unbalanced cycles, and so cannot  
 864 occur if  $(G, \mathcal{B})$  is tangled.  $\square$

## 865 4 Representations of unavoidable minors

866 In this section we examine the relationship between gain functions on the  
 867 biased graphs in  $\mathcal{G}_0$  (along with few other small biased graphs) and matrix  
 868 representations of their associated frame and lift matroids. In Section 4.1  
 869 we show that for each biased graph  $\Omega \in \mathcal{G}_0$ , a pair of canonical represen-  
 870 tations of  $F(\Omega)$  (resp.  $L(\Omega)$ ) are projectively equivalent if and only if their  
 871 associated gain functions are switching equivalent (resp. switching-and-scaling  
 872 equivalent). In Section 4.2 we show that for each biased graph  $\Omega \in \mathcal{G}_0$  ev-  
 873 ery  $\mathbb{F}$ -representation of each of  $F(\Omega)$  and  $L(\Omega)$  is projectively equivalent to a  
 874 canonical  $\mathbb{F}$ -representation particular to  $\Omega$ .

875 Our main tool for showing projective equivalence of a pair of matrix rep-  
 876 resentations is the following well-known result of Brylawski and Lucas.

877 **Proposition 4.1** ([10], Proposition 6.3.12). *Let  $A$  and  $B$  be  $r \times n$  matrices*  
 878 *over a field with the columns of each labelled, in order, by  $e_1, e_2, \dots, e_n$ ,*  
 879 *with  $r \geq 1$ . Then  $A$  and  $B$  are projectively equivalent representations of a*  
 880 *matroid on  $\{e_1, e_2, \dots, e_n\}$  if and only if there is a non-singular matrix  $T$  and*  
 881 *a non-singular diagonal matrix  $S$  such that  $TAS = B$ .*

## 882 4.1 Switching and projective equivalence

883 Let  $G$  be a graph and let  $\mathbb{F}$  be a field. Recall that for an  $\mathbb{F}^\times$ -gain function  $\varphi$   
 884 and an  $\mathbb{F}^+$ -gain function  $\psi$ , we denote by  $A_F(G, \varphi)$  and  $A_L(G, \psi)$ , respectively,  
 885 the canonical frame and lift matrices defined by  $(G, \varphi)$  and  $(G, \psi)$ , resp., as  
 886 described in Section 1.1. Our starting point is the following result of Zaslavsky.

887 **Proposition 4.2** (Zaslavsky [18]). *Let  $G$  be a graph and let  $\mathbb{F}$  be a field. Let*  
 888  *$\varphi$  and  $\psi$  be  $\mathbb{F}^\times$ - (resp.  $\mathbb{F}^+$ -)gain functions on  $G$ . If  $\varphi$  and  $\psi$  are switching*  
 889 *equivalent (resp. switching-and-scaling equivalent) then their canonical matrix*  
 890 *representations are projectively equivalent.*

891 *Proof.* Suppose  $\varphi$  and  $\psi$  are  $\mathbb{F}^\times$ -gain functions on  $G$  and  $\eta$  is a switching  
 892 function with  $\varphi^\eta = \psi$ . Let  $V(G) = \{v_1, v_2, \dots, v_{|V(G)|}\}$ . Let  $T$  be the diagonal  
 893 matrix with rows and columns indexed by  $V(G)$  in which diagonal entry  $T_{ii}$  is  
 894  $\eta(v_i)$ , and let  $S$  be the  $|E(G)| \times |E(G)|$  diagonal matrix with diagonal entries  
 895  $S_{jj} = \eta(v_i)^{-1}$  if vertex  $v_i$  is the tail of edge  $e_j$ . Then  $TA_F(G, \varphi)S = A_F(G, \psi)$ .


896 Now suppose  $\varphi$  and  $\psi$  are  $\mathbb{F}^+$ -gain functions,  $\eta$  is a switching function,  
 897 and that there is a scalar  $s \in \mathbb{F}^\times$  so that  $s\varphi^\eta = \psi$ . Let  $T$  be the  $(n+1) \times$   
 898  $(n+1)$  matrix whose first row is  $(s \ s\eta(v_1) \ s\eta(v_2) \ \dots \ s\eta(v_n))$ , first column is  
 899  $(s \ 0 \ 0 \ \dots \ 0)^T$ , and with the  $n \times n$  identity matrix as the submatrix consisting of  
 900 its remaining rows and columns. Let  $S$  be the diagonal matrix with  $s_{11} = 1/s$   
 901 and all other  $s_{ii} = 1$ . Then  $TA_L(G, \varphi)S = A_L(G, \psi)$ .  $\square$

902 The proof of Proposition 4.2 shows that if  $\varphi$  and  $\psi$  are switching equivalent  
 903  $\mathbb{F}^\times$ -gain functions, then a diagonal matrix  $T$  provides witness to the projective  
 904 equivalence of  $A_F(G, \varphi)$  and  $A_F(G, \psi)$ . The converse is also true. A similar  
 905 statement holds for canonical lift matrices.

906 **Lemma 4.3.** *Let  $G$  be a loopless graph and let  $\mathbb{F}$  be a field.*

907 1. *Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^\times$ -gain functions on  $G$  and assume  $A_F(G, \varphi)$  and*  
 908  *$A_F(G, \psi)$  are projectively equivalent. Then  $\varphi$  and  $\psi$  are switching equiv-*  
 909 *alent if and only if there exists a nonsingular diagonal matrix  $T$  and a*  
 910 *diagonal column-scaling matrix  $S$  such that  $TA_F(G, \varphi)S = A_F(G, \psi)$ .*

911 2. *Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^+$ -gain functions on  $G$  and assume  $A_L(G, \varphi)$  and  $A_L(G, \psi)$*   
 912 *are projectively equivalent. Then  $\varphi$  and  $\psi$  are switching-and-scaling*  
 913 *equivalent if and only if there exists a nonsingular matrix  $T$  and a diag-*  
 914 *onal column-scaling matrix  $S$  such that  $TA_L(G, \varphi)S = A_L(G, \psi)$ , where*  
 915 *removing the row and column of  $T$  indexed by  $v_0$  leaves an identity ma-*  
 916 *trix.*


Figure 9: Gain function  $\varphi$  on  $2C_3$ 

917 *Proof.* (1) Put  $A = A_F(G, \varphi)$  and  $B = A_F(G, \psi)$ . If  $\varphi$  and  $\psi$  are switching  
 918 equivalent, then the proof of Proposition 4.2 shows that  $A$  and  $B$  are pro-  
 919 jectively equivalent via diagonal nonsingular matrices  $T$  and  $S$ . Conversely,  
 920 suppose that  $B = TAS$  where  $T$  and  $S$  are both diagonal and nonsingular.  
 921 Since  $T$  is diagonal, row  $i$  of  $TA$  is obtained by multiplying row  $v_i$  of  $A$  by  
 922  $T_{ii}$ . Since both  $A$  and  $B$  are canonical frame representations, both have 1 in  
 923 position  $v_i$  of column  $e_j$  whenever vertex  $v_i$  is the tail of edge  $e_j$ . Hence the  
 924 diagonal elements  $S_{jj}$  of  $S$  satisfy  $S_{jj} = T_{ii}^{-1}$ , where  $v_i$  is the tail of  $e_j$ . Thus  
 925  $\varphi$  and  $\psi$  are switching equivalent via  $\eta(v_i) = T_{ii}$  for each  $v_i \in V(G)$ .

926 (2) Put  $A = A_L(G, \varphi)$  and  $B = A_L(G, \psi)$ . If  $\varphi$  and  $\psi$  are switching-and-  
 927 scaling equivalent, then the proof of Proposition 4.2 shows that  $A$  and  $B$  are  
 928 projectively equivalent via matrices  $T$  and  $S$  of the required forms. Conversely,  
 929 suppose that  $B = TAS$  where  $T$  and  $S$  are of the form in statement (2). Put  
 930  $n = |V(G)|$ . Index the rows and columns of  $T$  by  $\{v_0, v_1, \dots, v_n\}$ , where as  
 931 usual row  $v_0$  of  $A$  contains the gains assigned by  $\varphi$  and for  $i \in \{1, \dots, n\}$  row  
 932  $v_i$  corresponds to vertex  $v_i$ . Let  $\eta : V(G) \rightarrow \mathbb{F}^+$  be the switching function  
 933 defined by  $\eta(v_i) = T_{v_0 v_i}$  for  $i \in \{1, \dots, n\}$ . Then  $(T_{v_0 v_0}) \varphi^\eta = \psi$ .  $\square$

934 **Lemma 4.4.** Let  $\varphi : \vec{E}(2C_3) \rightarrow \mathbb{F}^\times$  be a gain function with  $\mathcal{B}_\varphi$  containing no  
 935 2-cycle, and let  $\psi : \vec{E}(2C_3) \rightarrow \mathbb{F}^\times$  be another gain function. Then  $\varphi$  and  $\psi$  are  
 936 switching equivalent if and only if  $A_F(2C_3, \varphi)$  and  $A_F(2C_3, \psi)$  are projectively  
 937 equivalent.

938 *Proof.* If  $\varphi$  and  $\psi$  are switching equivalent, then  $A_F(2C_3, \varphi)$  and  $A_F(2C_3, \psi)$   
 939 are projectively equivalent by Proposition 4.2. To prove the converse, let  
 940  $A = A_F(2C_3, \varphi)$  and  $B = A_F(2C_3, \psi)$  be a pair of projectively equivalent  
 941 canonical frame matrices. By normalizing on the spanning tree with edge set  
 942  $\{e_1, e_3\}$ , by Proposition 4.2 we may assume that  $\varphi$  assigns gains to  $E(2C_3)$  as

Figure 10: Gain function  $\varphi: \vec{E}(2C_3) \rightarrow \mathbb{F}^+$ .

943 shown in Figure 9. Then

$$944 \quad A = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 & 0 & -c & -d \\ -1 & -a & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -b & 1 & 1 \end{pmatrix} \end{matrix} \quad (4.1)$$

945 Since  $\mathcal{B}_\varphi$  contains no 2-cycle,  $a \neq 1 \neq b$  and  $c \neq d$ . Since  $A$  and  $B$  are  
 946 projectively equivalent, there is a nonsingular matrix  $T$  and a diagonal matrix  
 947  $S$  so that  $TA = BS$ . Denote by  $t_i$  row  $i$  of  $T$ , and by  $e_j$  column  $j$  of  $A$ .  
 948 Entry  $(BS)_{ij} = 0$  if and only if entry  $(TA)_{ij} = 0$ ; consider the dot products  
 949  $t_i \cdot e_j = 0$ , where  $(i, j) \in \{(3, 1), (3, 2), (1, 3), (1, 4), (2, 5), (2, 6)\}$ . The product  
 950  $t_3 \cdot e_1 = 0$  implies  $T_{31} = T_{32}$ , and  $t_3 \cdot e_2 = 0$  implies  $T_{31} = aT_{32}$ . Together  
 951 these imply (since  $a \neq 1$ ) that  $T_{32} = T_{31} = 0$ . Similarly,  $t_1 \cdot e_3 = t_1 \cdot e_4 = 0$   
 952 imply  $T_{13} = T_{12} = 0$ , and  $t_2 \cdot e_5 = t_2 \cdot e_6 = 0$  imply  $T_{21} = T_{23} = 0$ . Hence  $T$  is  
 953 diagonal, and so by Lemma 4.3,  $\varphi$  and  $\psi$  are switching equivalent.  $\square$

954 **Lemma 4.5.** *Let  $\varphi: \vec{E}(2C_3) \rightarrow \mathbb{F}^+$  be a gain function with  $\mathcal{B}_\varphi$  containing no*  
 955 *2-cycle and let  $\psi: \vec{E}(2C_3) \rightarrow \mathbb{F}^+$  be another gain function. Then  $\varphi$  and  $\psi$  are*  
 956 *switching-and-scaling equivalent if and only if  $A_L(2C_3, \varphi)$  and  $A_L(2C_3, \psi)$  are*  
 957 *projectively equivalent.*

958 *Proof.* If  $\varphi$  and  $\psi$  are switching-and-scaling equivalent, then their associated  
 959 lift matrices are projectively equivalent by Proposition 4.2. Conversely, let  
 960  $A = A_L(2C_3, \varphi)$  and  $B = A_L(2C_3, \psi)$  be a pair of projectively equivalent  
 961 canonical lift matrices. Normalizing on spanning tree  $\{e_1, e_2\}$ , and scaling if  
 962 necessary, we may assume that  $\varphi$  assigns gains to  $2C_3$  as shown in Figure 10.  
 963 Then

$$964 \quad A = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{matrix} v_0 \\ v_1 \\ v_2 \\ v_3 \end{matrix} & \begin{pmatrix} 0 & 1 & 0 & a & b & c \\ 1 & 1 & 0 & 0 & -1 & -1 \\ -1 & -1 & 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & -1 & 1 & 1 \end{pmatrix} \end{matrix} \quad (4.2)$$

965 Since  $\varphi$  has no balanced cycles of length 2,  $a \neq 0$  and  $b \neq c$ . There is a  
 966 non-singular matrix  $T$  and a diagonal matrix  $S$  so that  $TA = BS$ . As with  
 967  $\varphi$ , by switching and scaling we may assume  $\psi$  assigns gains to  $\vec{E}(2C_3)$  as in  
 968 Figure 10, replacing  $a$ ,  $b$ , and  $c$  with  $x$ ,  $y$ , and  $z$ , respectively. Then, denoting  
 969 elements  $S_{ii}$  of  $S$  by  $s_i$ , we have

$$970 \quad BS = \begin{pmatrix} 0 & s_2 & 0 & s_4x & s_5y & s_6z \\ s_1 & s_2 & 0 & 0 & -s_5 & -s_6 \\ -s_1 & -s_2 & s_3 & s_4 & 0 & 0 \\ 0 & 0 & -s_3 & -s_4 & s_5 & s_6 \end{pmatrix}$$

971 This gives 24 relations among the members of  $T$ , one for each dot product  
 972  $t_i \cdot e_j$ , where  $t_i$  is the  $i$ th column of  $T$  and  $e_j$  is the  $j$ th column of  $A$ . The eight  
 973 relations  $t_i \cdot e_j = 0$  yield  $T_{12} = T_{13} = T_{14}$ ,  $T_{21} = T_{31} = T_{41} = 0$ ,  $T_{23} = T_{24}$ , and  
 974  $T_{42} = T_{43}$ . Now after establishing these relations,  $t_3 \cdot e_5 = 0$  yields  $T_{32} = T_{34}$   
 975 and so

$$976 \quad T = \begin{pmatrix} T_{11} & T_{12} & T_{12} & T_{12} \\ 0 & T_{22} & T_{23} & T_{23} \\ 0 & T_{32} & T_{33} & T_{32} \\ 0 & T_{42} & T_{42} & T_{44} \end{pmatrix}$$

977 Now the relations  $s_1 = t_2 \cdot e_1$ ,  $s_2 = t_2 \cdot e_2$ ,  $s_3 = t_3 \cdot e_3$ ,  $s_4 = t_3 \cdot e_4$ ,  $s_5 = t_4 \cdot e_5$ ,  
 978  $s_6 = t_4 \cdot e_6$ ,  $-s_2 = t_3 \cdot e_2$ , and  $-s_3 = t_4 \cdot e_3$  yield  $s_1 = s_2 = s_3 = s_4 = s_5 = s_6$ .  
 979 Hence the relation  $s_2 = t_1 \cdot e_2$  yields  $T_{11} = s_1$ . Now the relations  $t_1 \cdot e_4 = s_1x$ ,  
 980  $t_1 \cdot e_5 = s_1y$ , and  $t_1 \cdot e_6 = s_1z$  yield  $a = x$ ,  $b = y$  and  $c = z$ . This implies that  
 981  $\varphi$  and  $\psi$  are switching equivalent after scaling.  $\square$

982 **Lemma 4.6.** *Let  $\varphi: \vec{E}(2C_3) \rightarrow \mathbb{F}^\times$  and  $\psi: \vec{E}(2C_3) \rightarrow \mathbb{F}^+$  be gain functions,*  
 983 *neither of which yield a balanced 2-cycle. Then  $A_F(2C_3, \varphi)$  and  $A_L(2C_3, \psi)$*   
 984 *are not projectively equivalent.*

985 *Proof.* As in previous cases, without loss of generality we may assume that  $\varphi$   
 986 assigns gains to  $2C_3$  as in Figure 9, and  $\psi$  as in Figure 10, replacing  $a$  with  $x$ ,  $b$   
 987 with  $y$ , and  $c$  with  $z$ . Then  $A_F(2C_3, \varphi)$  is the matrix of (4.1), and  $A_L(2C_3, \psi)$   
 988 is the matrix of (4.2) with  $a$ ,  $b$ , and  $c$  replaced by  $x$ ,  $y$ , and  $z$  respectively. Let  
 989  $B$  be the matrix obtained from  $A_L(2C_2, \psi)$  by removing row  $v_3$ ; then  $B$  is a  
 990 canonical lift matrix of full rank given by  $\psi$ . Now suppose for a contradiction  
 991 that there exists a non-singular matrix  $T$  and a diagonal matrix  $S$  so that  
 992  $TA = BS$ . Writing  $S_{ii} = s_i$ , and denoting row  $i$  of  $T$  by  $t_i$  and column  $j$  of  $A$   
 993 by  $e_j$ , we have  $t_2 \cdot e_1 = s_1$ ,  $t_2 \cdot e_2 = s_2$ ,  $t_2 \cdot e_3 = 0$ , and  $t_2 \cdot e_4 = 0$ . Together  
 994 these imply that  $T_{22} = T_{23} = 0$  and that  $T_{21} = s_1 = s_2$ . Moreover, we have  
 995  $t_3 \cdot e_1 = -s_1$ ,  $t_3 \cdot e_2 = -s_2$ ,  $t_3 \cdot e_5 = 0$ , and  $t_3 \cdot e_6 = 0$ . Since  $s_1 = s_2$ ,  $a \neq 0, 1$ ,

996 and  $c \neq d$ , these imply that  $T_{31} = T_{32} = T_{33} = 0$  which implies that  $T$  is  
 997 singular, a contradiction.  $\square$

998 **Lemma 4.7.** *Let  $\varphi: \vec{E}(K_4) \rightarrow \mathbb{F}^\times$  be a gain function with  $\mathcal{B}_\varphi$  containing no  
 999 3-cycle, and let  $\psi: \vec{E}(K_4) \rightarrow \mathbb{F}^\times$  be another gain function. Then  $\varphi$  and  $\psi$  are  
 1000 switching equivalent if and only if  $A_F(K_4, \varphi)$  and  $A_F(K_4, \psi)$  are projectively  
 1001 equivalent.*

1002 *Proof.* By switching we may assume that  $\varphi$  and  $\psi$  are both equal to the iden-  
 1003 tity on a  $K_{1,3}$ -subgraph  $Y$ . This allows us to consider  $\varphi$  and  $\psi$  as gain functions  
 1004 on  $\nabla_Y K_4 = 2C_3$ . Now  $\nabla_Y(K_4, \mathcal{B}_\varphi) = (2C_3, \mathcal{B}_\varphi)$  and  $\nabla_Y(K_4, \mathcal{B}_\psi) = (2C_3, \mathcal{B}_\psi)$ .  
 1005 Since  $\varphi$  has no balanced triangles in  $K_4$ , neither has it any balanced 2-cycles  
 1006 in  $2C_3$ . So Lemma 4.4 implies that  $\varphi$  and  $\psi$  are switching equivalent if and  
 1007 only if  $A_F(2C_3, \varphi)$  and  $A_F(2C_3, \psi)$  are projectively equivalent and so Propo-  
 1008 sitions 2.10 and 2.9 imply that  $\varphi$  and  $\psi$  are switching equivalent if and only if  
 1009  $A_F(K_4, \varphi)$  and  $A_F(K_4, \psi)$  are projectively equivalent.  $\square$

1010 Using  $Y$ - $\Delta$  exchanges as in the proof of Lemma 4.7 yields Lemmas 4.8 and  
 1011 4.9.


1012 **Lemma 4.8.** *Let  $\varphi: \vec{E}(K_4) \rightarrow \mathbb{F}^+$  be a gain function with  $\mathcal{B}_\varphi$  containing no  
 1013 3-cycle and let  $\psi: \vec{E}(K_4) \rightarrow \mathbb{F}^+$  be another gain function. Then  $\varphi$  and  $\psi$  are  
 1014 switching-and-scalaing equivalent if and only if  $A_L(K_4, \varphi)$  and  $A_L(K_4, \psi)$  are  
 1015 projectively equivalent.*

1016 **Lemma 4.9.** *Let  $\varphi: \vec{E}(K_4) \rightarrow \mathbb{F}^\times$  and  $\psi: \vec{E}(K_4) \rightarrow \mathbb{F}^+$  be gain functions  
 1017 neither of which yield a balanced 3-cycle. Then  $A_F(K_4, \varphi)$  and  $A_L(K_4, \psi)$  are  
 1018 not projectively equivalent.*

1019 **Lemma 4.10.** *Let  $\varphi: \vec{E}(2C_4'') \rightarrow \mathbb{F}^\times$  be a gain function with  $\mathcal{B}_\varphi$  containing no  
 1020 2-cycle, and let  $\psi: \vec{E}(2C_4'') \rightarrow \mathbb{F}^\times$  be another gain function. Then  $\varphi$  and  $\psi$  are  
 1021 switching equivalent if and only if  $A_F(2C_4'', \varphi)$  and  $A_F(2C_4'', \psi)$  are projectively  
 1022 equivalent.*

1023 *Proof.* Let  $A = A_F(2C_4'', \varphi)$  and  $B = A_F(2C_4'', \psi)$ . If  $\varphi$  and  $\psi$  are switching  
 1024 equivalent, then  $A$  and  $B$  are projectively equivalent by Proposition 4.2. To  
 1025 prove the converse, let  $T$  and  $S$  be matrices with  $TA = BS$ , where  $T$  is  
 1026 nonsingular and  $S$  is a diagonal matrix scaling the columns of  $B$ . We may  
 1027 assume without loss of generality that the edge orientations chosen to define  
 1028  $B$  are the same as those chosen to define  $A$ ; by normalizing on the spanning



Figure 11: Labeled  $2C_4''$  with a normalized gain function.

1029 tree with edges  $e_3, e_4, e_5$ , we may assume without loss of generality that  $\varphi$   
 1030 assigns gains to  $\vec{E}(K_4)$  as shown in Figure 11 where  $a \neq b$  and  $c \neq 1$ . Thus

$$1031 \quad A = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{matrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{matrix} & \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ -a & -b & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & 1 \\ 0 & 0 & 0 & -1 & -1 & -c \end{pmatrix} \end{matrix}.$$

1032 Now, each entry  $B_{ij}$  of  $B$  is zero if and only if entry  $(TA)_{ij} = 0$ . Now for a  
 1033 fixed row  $i$ , there are three distinct  $j$  such that  $t_i \cdot e_j = 0$ . It is a straightforward  
 1034 check that for each  $i$ , that these three relations yield  $T_{ij} \neq 0$  if and only if  
 1035  $i = j$ . For  $i = 1$ , the relations are  $0 = t_1 \cdot e_4$ , which implies  $T_{12} = T_{14}$ ;  
 1036  $0 = t_1 \cdot e_5$ , which implies  $T_{13} = T_{14}$ ; and  $0 = t_1 \cdot e_6$ , which implies  $T_{13} = cT_{14}$ .  
 1037 Since  $c \neq 1$ , this implies  $T_{12} = T_{13} = T_{14} = 0$ . Similarly, the entries of  $T$  off its  
 1038 main diagonal in rows 2, 3, and 4 are all zero. Thus  $T$  is diagonal. By Lemma  
 1039 4.3,  $\varphi$  and  $\psi$  are switching equivalent.  $\square$

1040 **Lemma 4.11.** *Let  $\varphi: \vec{E}(2C_4'') \rightarrow \mathbb{F}^+$  be a gain function with  $\mathcal{B}_\varphi$  containing*  
 1041 *no 2-cycle, and let  $\psi: \vec{E}(2C_4'') \rightarrow \mathbb{F}^+$  be another gain function. Then  $\varphi$  and  $\psi$*   
 1042 *are switching-and-scaling equivalent if and only if  $A_L(2C_4'', \varphi)$  and  $A_L(2C_4'', \psi)$*   
 1043 *are projectively equivalent.*

1044 *Proof.* The “only if” statement again follows from Proposition 4.2. For the  
 1045 converse, without loss of generality assume that  $\varphi(e_1) = \psi(e_1) = 1$ ,  $\varphi(e_2) = a$ ,  
 1046  $\psi(e_2) = x$ ,  $\varphi(e_3) = \psi(e_3) = 0$ ,  $\varphi(e_4) = \psi(e_4) = 0$ ,  $\varphi(e_5) = \psi(e_5) = 0$ ,  
 1047  $\varphi(e_6) = b$ , and  $\psi(e_6) = y$  (where  $2C_4''$  has edges and orientations as in Figure  
 1048 11) such that neither  $a$  nor  $x$  is 1 and neither  $b$  nor  $y$  is 0. Let  $A = A_L(2C_4'', \varphi)$   
 1049 and  $B = A_L(2C_4'', \psi)$ , and let  $T$  and  $S$  be matrices with  $TA = BS$ , where  $S$  is  
 1050 diagonal (with  $s_i = S_{ii}$ ) scaling the columns of  $B$ . Denoting row  $i$  of  $T$  by  $t_i$   
 1051 and column  $j$  of  $A$  by  $e_j$  we have  $t_i \cdot e_j = 0$  for 15 pairs  $(i, j)$ , three pairs for




Figure 12

1052 each row. It is straightforward to deduce that these relations imply

$$1053 \quad T = \begin{pmatrix} T_{11} & T_{12} & T_{12} & T_{12} & T_{12} \\ 0 & T_{22} & T_{23} & T_{23} & T_{23} \\ 0 & T_{32} & T_{33} & T_{32} & T_{32} \\ 0 & T_{42} & T_{42} & T_{44} & T_{42} \\ 0 & T_{52} & T_{52} & T_{52} & T_{55} \end{pmatrix}.$$

1054 Now each column  $e_j$  has  $i, k \geq 2$  such that  $t_i \cdot e_j = s_j$  and  $t_k \cdot e_j = -s_j$ . These  
 1055 12 relations yield  $s_1 = s_2 = s_3 = s_4 = s_5 = s_6$ . The relation  $t_1 \cdot e_1 = s_1$  yields  
 1056  $T_{11} = s_1$ . Now the relation  $t_1 \cdot e_2 = s_1 x$  yields  $a = x$  and the relation  $t_1 \cdot e_6 = s_1 y$   
 1057 yields  $b = y$ . Thus  $A$  and  $B$  are switching-and-scaling equivalent.  $\square$

1058 Two biased graphic representations of  $U_{2,4}$  are  $U_2$  and  $U_3$ , shown in Figure  
 1059 7, where all cycles in each are unbalanced. Denote the underlying graphs of  
 1060  $U_2$  and  $U_3$  by  $U_2$  and  $U_3$ , respectively.

1061 **Lemma 4.12.** *Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^\times$ -realizations of  $U_2$ . Then  $A_F(U_2, \varphi)$  and*  
 1062  *$A_F(U_2, \psi)$  are projectively equivalent if and only if  $\varphi(e_3 e_4) = \psi(e_3 e_4)$ .*

*Proof.* We may assume  $U_2$  is labeled with edge orientations as in Figure 12. Matrices  $A_F(U_2, \varphi)$  and  $A_F(U_2, \psi)$  are of the form

$$\begin{pmatrix} 1 & 0 & 1 & -g \\ 0 & 1 & -1 & 1 \end{pmatrix} \tag{4.3}$$

1063 up to scaling columns  $e_1$  and  $e_2$ . These are in standard form relative to the  
 1064 basis  $\{e_1, e_2\}$  and so are projectively equivalent if and only if entry  $g$  is the  
 1065 same for both  $A_F(U_2, \varphi)$  and  $A_F(U_2, \psi)$ . The result follows.  $\square$

1066 **Lemma 4.13.** *Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^+$ -realizations of  $U_3$ . Then  $A_L(U_3, \varphi)$  and*  
 1067  *$A_L(U_3, \psi)$  are projectively equivalent if and only if  $\varphi|_{\{e_2, e_3, e_4\}}$  and  $\psi|_{\{e_2, e_3, e_4\}}$*   
 1068 *are switching-and-scaling equivalent.*

1069 *Proof.* We may assume that  $U_3$  is labelled as in Figure 12. If  $A = A_L(U_3, \varphi)$   
 1070 and  $B = A_L(U_3, \psi)$  are projectively equivalent, then there is an invertible  
 1071 matrix  $T$  and diagonal matrix  $S$  such that  $TA = BS$ . By switching and scaling  
 1072 we may assume that  $\varphi(e_1) = \psi(e_1) = 1$ ,  $\varphi(e_2) = \psi(e_2) = 0$ ,  $\varphi(e_3) = \psi(e_3) = 1$ ,  
 1073  $\varphi(e_4) = a$ , and  $\psi(e_4) = x$ . Again writing  $s_i$  for entry  $S_{ii}$ ,

$$1074 \begin{pmatrix} T_{11} & T_{12} & T_{13} \\ T_{21} & T_{22} & T_{23} \\ T_{31} & T_{32} & T_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 & a \\ 0 & 1 & 1 & -1 \\ 0 & -1 & -1 & 1 \end{pmatrix} = \begin{pmatrix} s_1 & 0 & s_3 & s_4x \\ 0 & s_2 & s_3 & -s_4 \\ 0 & -s_2 & -s_3 & s_4 \end{pmatrix}.$$

1075 This yields  $T_{11} = s_1$ ,  $T_{21} = T_{31} = 0$ , and  $T_{12} = T_{13}$ . That is,

$$1076 \begin{pmatrix} s_1 & T_{12} & T_{12} \\ 0 & T_{22} & T_{23} \\ 0 & T_{32} & T_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 & a \\ 0 & 1 & 1 & -1 \\ 0 & -1 & -1 & 1 \end{pmatrix} = \begin{pmatrix} s_1 & 0 & s_3 & s_4x \\ 0 & s_2 & s_3 & -s_4 \\ 0 & -s_2 & -s_3 & s_4 \end{pmatrix},$$

1077 which yields  $s_1 = s_2 = s_3 = s_4$ . Thus  $a = x$  and so  $\varphi$  and  $\psi$  are switching-  
 1078 and-scaling equivalent.  $\square$

## 1079 4.2 All $\mathbb{F}$ -representations are canonical

1080 Let  $\mathbb{F}$  be a field. In this section we show that every  $\mathbb{F}$ -matrix representation of a  
 1081 frame or lift matroid arising from a biased graph in  $\mathcal{T}_0$  is projectively equivalent  
 1082 to a canonical representation particular to that biased graph. Recall that  
 1083 when  $(G, \mathcal{B})$  is a biased graph with no two vertex-disjoint unbalanced cycles,  
 1084  $F(G, \mathcal{B}) = L(G, \mathcal{B})$ , and we denote this common matroid by  $M(G, \mathcal{B})$ .

1085 **Lemma 4.14.** *Let  $(2C_3, \mathcal{B})$  be a biased graph with no balanced 2-cycle and let*  
 1086  *$A$  be an  $\mathbb{F}$ -matrix representing  $M(2C_3, \mathcal{B})$ . Then  $A$  is projectively equivalent*  
 1087 *to a canonical lift matrix particular to  $(2C_3, \mathcal{B})$  or to a canonical frame matrix*  
 1088 *particular to  $(2C_3, \mathcal{B})$ , but not both.*

1089 *Proof.* We may assume that  $2C_3$  is labeled and has edge orientations as shown  
 1090 in Figure 13. Let  $A$  be a matrix over  $\mathbb{F}$  representing  $M(2C_3, \mathcal{B})$ . If  $(2C_3, \mathcal{B}) \cong$   
 1091  $\mathsf{T}_4$  then  $M(2C_3, \mathcal{B})$  is isomorphic to the cycle matroid of  $K_4$ , and so has a  
 1092 projectively unique representation over every field. Thus if the characteristic  
 1093 of  $\mathbb{F}$  is two then  $A$  is projectively equivalent to the canonical lift matrix

$$1094 B = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 & 1 \end{pmatrix}$$




Figure 13

1095 and if the characteristic of  $\mathbb{F}$  is not two then  $A$  is projectively equivalent to  
 1096 the canonical frame matrix

$$1097 \quad C = \begin{pmatrix} 1 & 0 & 1 & 1 & 0 & -1 \\ 1 & 1 & 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 0 & -1 & 1 \end{pmatrix}.$$

1098 We now claim that there is no canonical frame matrix particular to  $\mathbb{T}_4$  over  
 1099 any field of characteristic two, and that neither is there a canonical lift matrix  
 1100 particular to  $\mathbb{T}_4$  in any field of characteristic different from two. For, toward  
 1101 a contradiction, suppose  $D$  is a canonical frame matrix particular to  $\mathbb{T}_4$  over  
 1102 a field  $\mathbb{F}$  of characteristic two. Assume the collection of balanced cycles of  $\mathbb{T}_4$   
 1103 is  $\{e_1e_2e_6, e_1e_3e_5, e_2e_3e_4, e_4e_5e_6\}$ . We may assume

$$1104 \quad D = \begin{pmatrix} e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ 1 & 0 & c & 1 & 0 & 1 \\ a & 1 & 0 & 1 & 1 & 0 \\ 0 & b & 1 & 0 & 1 & 1 \end{pmatrix}$$

1105 where  $a, b, c \in \mathbb{F}^\times$  (and we omit the customary negative signs as redundant).  
 1106 Since  $e_1e_2e_6$  is balanced,  $ab = 1$ ; since  $e_1e_3e_5$  is balanced,  $ac = 1$ ; and because  
 1107  $e_2e_3e_4$  is balanced,  $bc = 1$ . These relations imply that  $a = b = c$  and so  
 1108 that  $a^2 = 1$ . But this implies  $a = 1$ , and so  $D$  does not represent  $M(\mathbb{T}_4)$ , a  
 1109 contradiction.

1110 Similarly, suppose for a contradiction that  $D$  is a canonical lift matrix  
 1111 particular to  $\mathbb{T}_4$  over a field  $\mathbb{F}$  of characteristic different from two. Then we  
 1112 may assume

$$1113 \quad D = \begin{pmatrix} e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ 0 & 0 & 1 & a & b & c \\ 1 & 0 & -1 & 1 & 0 & -1 \\ -1 & 1 & 0 & -1 & 1 & 0 \end{pmatrix}$$

1114 for some nonzero elements  $a, b, c \in \mathbb{F}$ , where the second and third rows are the  
 1115 oriented incidence matrix of  $2C_3$  with its row corresponding to  $v_3$  removed.  
 1116 Because  $e_4e_5e_6$  is balanced,  $a + b + c = 0$ . Since  $e_1e_2e_6$  is balanced,  $c = 0$ ;  
 1117 since  $e_1e_3e_5$  is balanced,  $1 + b = 0$ ; and because  $e_2e_3e_4$  is balanced,  $1 + a = 0$ .  
 1118 These relations imply that  $a = b = -1$  and so that  $a + b + c = -2 \neq 0$ , a  
 1119 contradiction. This completes the proof in the case that  $(2C_3, \mathcal{B}) \cong T_4$ .

1120 Now assume  $(2C_3, \mathcal{B}) \not\cong T_4$ . By Proposition 3.1 we may assume that the  
 1121 triangles  $\{e_1, e_2, e_3\}$  and  $\{e_2, e_3, e_4\}$  are both unbalanced. Since the only form  
 1122 a 3-circuit takes in  $(2C_3, \mathcal{B})$  is a balanced triangle and neither  $e_5$  nor  $e_6$  forms  
 1123 a triangle with  $\{e_2, e_3\}$ ,  $A$  is projectively equivalent to the matrix

$$1124 \quad \begin{array}{cccccc} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ & \left( \begin{array}{cccccc} 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & a & c \\ 0 & 0 & 1 & 1 & b & d \end{array} \right). \end{array}$$

1125 Hence:

- 1126 1. Neither  $b$  nor  $c$  is 0. If  $b = 0$  then  $\{e_1, e_2, e_5\}$  is a circuit; if  $c = 0$  then  
 1127  $\{e_1, e_3, e_6\}$  is a circuit: both contradictions.
- 1128 2.  $a \neq b$ : If so then  $a \neq 1$ , as then  $e_4$  and  $e_5$  would form a parallel pair, a  
 1129 contradiction. But then  $\{e_1, e_4, e_5\}$  is a circuit, also a contradiction.
- 1130 3.  $b \neq 1$ : If so, then  $\{e_2, e_4, e_5\}$  is a circuit, a contradiction.
- 1131 4.  $c \neq d$ : If so, then  $c \neq 1$  as  $e_4$  and  $e_6$  are not a parallel pair. But then  
 1132  $\{e_1, e_4, e_6\}$  is a circuit, a contradiction.
- 1133 5.  $c \neq 1$ : If so,  $\{e_3, e_4, e_6\}$  is a circuit, a contradiction.
- 1134 6.  $a \neq c$ : If so, then  $d \neq b$  since  $e_5$  and  $e_6$  are not a parallel pair. But then  
 1135  $\{e_3, e_5, e_6\}$  is a circuit, a contradiction.
- 1136 7.  $b \neq d$ : If so, then  $a \neq c$  since  $e_5$  and  $e_6$  are not a parallel pair. But then  
 1137  $\{e_2, e_5, e_6\}$  is a circuit, a contradiction.

1138 Suppose there are nonsingular matrices  $T$  and  $S$  such that  $TAS$  is a canon-  
 1139 ical frame matrix particular to  $(2C_3, \mathcal{B})$ , where  $S$  is diagonal column-scaling  
 1140 matrix. Then we may assume

$$1141 \quad TAS = \begin{array}{cccccc} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ & \left( \begin{array}{cccccc} 1 & 0 & -g_1 & 1 & 0 & -g_4 \\ -1 & 1 & 0 & -g_2 & 1 & 0 \\ 0 & -1 & 1 & 0 & -g_3 & 1 \end{array} \right) \end{array}$$

1142 for some elements  $g_1, \dots, g_4 \in \mathbb{F} - \{0, 1\}$ . Let us denote row  $i$  of  $T$  by  $t_i$  and  
 1143 column  $j$  of  $A$  by  $e_j$ . Consider the products  $t_i \cdot e_j = (TA)_{ij}$  for  $1 \leq i \leq 3$ ,  
 1144  $1 \leq j \leq 6$ . The products  $t_3 \cdot e_1 = 0$ ,  $t_1 \cdot e_2 = 0$ , and  $t_2 \cdot e_3 = 0$  imply  $T_{31} = 0$ ,  
 1145  $T_{12} = 0$ , and  $T_{23} = 0$ , respectively. Since  $t_1 \cdot e_1 = -(t_2 \cdot e_1)$ ,  $T_{21} = -T_{11}$ .  
 1146 Similarly,  $t_2 \cdot e_2 = -(t_3 \cdot e_2)$  implies  $T_{32} = -T_{22}$ . Now  $t_3 \cdot e_4 = 0$  implies  
 1147  $T_{33} = -T_{32}$ ;  $t_1 \cdot e_5 = 0$  implies  $T_{13} = -T_{11}/b$ ; and finally,  $t_2 \cdot e_6 = 0$  implies  
 1148 that  $T_{22} = -T_{21}/c$ . Thus  $T$  is the matrix

$$1149 \quad \begin{pmatrix} t & 0 & -t/b \\ -t & t/c & 0 \\ 0 & -t/c & t/c \end{pmatrix} = t \begin{pmatrix} 1 & 0 & -1/b \\ -1 & 1/c & 0 \\ 0 & -1/c & 1/c \end{pmatrix}$$

1150 for some nonzero  $t \in \mathbb{F}$ . Since  $T$  has determinant  $t^3(1/c^2 - 1/bc)$ ,  $T$  is non-  
 1151 singular if and only if  $b \neq c$ . Assuming  $b \neq c$ , and taking  $t = 1$ ,

$$1152 \quad TA = \begin{pmatrix} 1 & 0 & -1/b & (b-1)/b & 0 & (b-d)/b \\ -1 & 1/c & 0 & (1-c)/c & (a-c)/c & 0 \\ 0 & -1/c & 1/c & 0 & (b-a)/c & (d-c)/c \end{pmatrix}.$$

1153 By claims 1-7 above  $TA$  has exactly two nonzero entries in each column, so  
 1154 scaling the columns of  $TA$  appropriately yields a canonical frame matrix. Thus  
 1155  $A$  is projectively equivalent to a canonical frame matrix particular to  $(2C_3, \mathcal{B})$   
 1156 if and only if  $b \neq c$ .

1157 Now let  $T$  be a nonsingular matrix such that  $TAS$  is a canonical lift matrix  
 1158 particular to  $(2C_3, \mathcal{B})$ , for some diagonal column-scaling matrix  $S$ . We may  
 1159 assume that  $TAS$  is of the form

$$1160 \quad TAS = \begin{pmatrix} 0 & 0 & 1 & g_1 & g_2 & g_3 \\ 1 & 0 & -1 & 1 & 0 & -1 \\ -1 & 1 & 0 & -1 & 1 & 0 \end{pmatrix}$$

1161 for some nonzero  $g_1, g_2, g_3 \in \mathbb{F}^+$ , where row 1 is indexed by  $v_0$  and rows 2 and  
 1162 3 are the oriented incidence matrix of  $2C_3$  with its row corresponding to  $v_3$   
 1163 removed. Consider the products  $t_i \cdot e_j = (TA)_{ij}$ . The products  $t_1 \cdot e_1 = 0$ ,  
 1164  $t_1 \cdot e_2 = 0$ ,  $t_2 \cdot e_2 = 0$ , and  $t_3 \cdot e_3 = 0$  imply  $T_{11} = 0$ ,  $T_{12} = 0$ ,  $T_{22} = 0$ , and  $T_{33} = 0$ ,  
 1165 respectively. Thus  $t_2 \cdot e_1 = -(t_3 \cdot e_1)$  implies  $T_{21} = -T_{31}$  and  $t_1 \cdot e_3 = -(t_2 \cdot e_3)$   
 1166 implies  $T_{13} = -T_{23}$ ;  $t_2 \cdot e_5 = 0$  implies  $T_{21} = -bT_{23}$  and  $t_3 \cdot e_6 = 0$  implies  
 1167  $T_{31} = -cT_{32}$ . Now  $t_2 \cdot e_4 = -(t_3 \cdot e_4)$  yields  $T_{23} = -T_{32}$ , which the preceding  
 1168 relations imply is equivalent to the statement  $T_{31}/b = T_{31}/c$ . This holds if and  
 1169 only if either  $T_{31} = 0$  or  $b = c$ . Hence if  $b \neq c$ ,  $T_{31} = 0$ . Then the preceding  
 1170 relations imply that  $T_{32} = 0$ . Since  $T_{33} = 0$ , this implies  $T$  is singular, a

1171 contradiction. Thus in the case  $b \neq c$ ,  $A$  is not projectively equivalent to a  
 1172 canonical lift matrix particular to  $(2C_3, \mathcal{B})$ .

1173 So assume  $T_{31}$  is nonzero and  $b = c$ . Then the relations above imply  $T$  is  
 1174 the matrix

$$1175 \quad \begin{pmatrix} 0 & 0 & t \\ bt & 0 & -t \\ -bt & t & 0 \end{pmatrix} = t \begin{pmatrix} 0 & 0 & 1 \\ b & 0 & -1 \\ -b & 1 & 0 \end{pmatrix}$$

1176 for some nonzero  $t \in \mathbb{F}$ . Matrix  $T$  is non-singular; taking  $t = 1$  we have

$$1177 \quad TA = \begin{pmatrix} 0 & 0 & 1 & 1 & b & d \\ -b & 0 & -1 & b-1 & 0 & b-d \\ b & 1 & 0 & 1-b & a-b & 0 \end{pmatrix}.$$

1178 By claims 1, 2, 3, and 7, none of  $b$ ,  $b-1$ ,  $a-b$ , nor  $b-d$  is zero. By scaling  
 1179 columns so that all nonzero entries in rows 2 and 3 are  $\pm 1$  (and appending a  
 1180 fourth row obtained by negating the sum of rows 2 and 3 if desired), we obtain  
 1181 a canonical lift matrix particular to  $(2C_3, \mathcal{B})$ . Thus  $A$  is projectively equivalent  
 1182 to a canonical lift matrix particular to  $(2C_3, \mathcal{B})$  if and only if  $b = c$ .  $\square$

1183 Recall that  $\mathbf{P}$  is the triangular prism with just its two triangles balanced,  
 1184 and that  $\mathbf{P}_1$  and  $\mathbf{P}_2$  are obtained from  $\mathbf{P}$  by contracting 2 and 1 of the edges  
 1185 of the matching between the two triangles, respectively (Figure 5).

1186 **Lemma 4.15.** *Let  $\Omega \in \{\mathbf{P}, \mathbf{P}_1, \mathbf{P}_2\}$  and let  $A$  be an  $\mathbb{F}$ -matrix representing  
 1187  $M(\Omega)$ . Then  $A$  is projectively equivalent to a canonical lift matrix particular  
 1188 to  $\Omega$  or to a canonical frame matrix particular to  $\Omega$ , but not both.*

1189 *Proof.* First, consider  $\mathbf{P}_1$ . Let  $A$  be an  $\mathbb{F}$ -representation of  $M(\mathbf{P}_1)$ . Let  $Y$  be  
 1190 a  $K_{1,3}$ -subgraph of  $\mathbf{P}_1$ . Then  $\nabla_Y \mathbf{P}_1$  is obtained from  $\mathbf{T}_2$  by the addition of an  
 1191 edge  $e$  that creates a balanced 2-cycle. Hence  $e$  is parallel with an element of  
 1192  $M(\nabla_Y \mathbf{P}_1)$ . By Proposition 2.8  $M(\nabla_Y \mathbf{P}_1) = \nabla_Y M(\mathbf{P}_1)$ . Thus by Lemma 4.14  
 1193 every  $\mathbb{F}$ -representation of  $\nabla_Y M(\mathbf{P}_1)$  is projectively equivalent to a canonical  
 1194 representation particular to  $\nabla_Y \mathbf{P}_1$ . In particular,  $\nabla_Y A$  is projectively equiv-  
 1195 alent to a canonical representation particular to  $\nabla_Y \mathbf{P}_1$ . Thus by Proposition  
 1196 2.11  $A$  is projectively equivalent to a canonical representation particular to  $\mathbf{P}_1$ .

1197 Now consider  $\mathbf{P}_2$ . Since  $\nabla_Y \mathbf{P}_2$  is obtained from  $\mathbf{P}_1$  by the addition of an  
 1198 edge that creates a balanced 2-cycle, by the argument analogous to that of the  
 1199 previous paragraph every  $\mathbb{F}$ -representation of  $M(\mathbf{P}_2)$  is projectively equivalent  
 1200 to a canonical representation particular to  $\mathbf{P}_2$ . Finally, the observation that  
 1201  $\nabla_Y \mathbf{P}$  is obtained from  $\mathbf{P}_2$  by the addition of an edge that creates a balanced  
 1202 2-cycle, along with the argument analogous to that of the previous paragraph,  
 1203 establishes the statement for  $M(\mathbf{P})$ .




Figure 14

1204 By Proposition 2.11 and Lemma 4.14, in no case may  $A$  be projectively  
 1205 equivalent to both a canonical lift and a canonical frame matrix.  $\square$

1206 **Lemma 4.16.** *Let  $(K_4, \mathcal{B})$  be a biased graph with no balanced 3-cycle and let*  
 1207  *$A$  be an  $\mathbb{F}$ -matrix representing  $M(K_4, \mathcal{B})$ . Then  $A$  is projectively equivalent*  
 1208 *to a canonical lift matrix particular to  $(K_4, \mathcal{B})$  or to a canonical frame matrix*  
 1209 *particular to  $(K_4, \mathcal{B})$ , but not both.*

1210 *Proof.* The biased graph  $(K_4, \mathcal{B})$  is isomorphic to  $D_{0,i}$  for some  $i \in \{0, 1, 2, 3\}$ .  
 1211 There is a  $K_{1,3}$ -subgraph  $Y$  of  $(K_4, \mathcal{B})$  so that  $\nabla_Y(K_4, \mathcal{B}) \cong T_{i+1}$ . Since  
 1212  $\nabla_Y M(K_4, \mathcal{B}) = M(\nabla_Y(K_4, \mathcal{B}))$ , the result follows by Lemma 4.14 and Propo-  
 1213 sition 2.11.  $\square$

1214 **Lemma 4.17.** *Let  $(2C_4'', \mathcal{B})$  be a biased graph with no balanced 2-cycle and let*  
 1215  *$A$  be an  $\mathbb{F}$ -matrix representing  $F(2C_4'', \mathcal{B})$ . Then  $A$  is projectively equivalent to*  
 1216 *a canonical frame matrix particular to  $(2C_4'', \mathcal{B})$ .*

1217 *Proof.* Without loss of generality we may assume that  $2C_4''$  is labelled as shown  
 1218 in Figure 14. There are three possibilities for  $\mathcal{B}$ :  $|\mathcal{B}| \in \{0, 1, 2\}$ .

1219 Assume first that  $|\mathcal{B}|$  is 0 or 1; *i.e.* either  $\mathcal{B} = \emptyset$  or, without loss of generality,  
 1220  $\mathcal{B} = \{e_1 e_3 e_4 e_6\}$ . Then  $A$  is projectively equivalent to the matrix

$$1221 \quad A' = \begin{pmatrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & a \\ 0 & 0 & 1 & 0 & 1 & b \\ 0 & 0 & 0 & 1 & 1 & c \end{pmatrix} \end{pmatrix}$$

1222 where  $a, b, c \in \mathbb{F}$  are distinct, neither of  $b$  nor  $c$  is 0, and none of  $a, b$ , or  $c$  are  
 1223 1; in the case that  $\mathcal{B} = \emptyset$ ,  $a \neq 0$ , while if  $\mathcal{B} = \{e_1 e_3 e_4 e_5\}$  then  $a = 0$ . Let

$$1224 \quad T = \begin{pmatrix} b-a & 1-b & a-1 & 0 \\ a-c & c-1 & 0 & 1-a \\ 0 & 0 & 1-a & 0 \\ 0 & 0 & 0 & a-1 \end{pmatrix}.$$



1225 Then  $\det(T) = (a - 1)^3(c - b)$  so  $T$  is nonsingular, and

$$1226 \quad TA' = \begin{pmatrix} b-a & 1-b & a-1 & 0 & 0 & 0 \\ a-c & c-1 & 0 & 1-a & 0 & 0 \\ 0 & 0 & 1-a & 0 & 1-a & (1-a)b \\ 0 & 0 & 0 & a-1 & a-1 & (a-1)c \end{pmatrix}$$

1227 which has the desired canonical form after column scaling.

1228 So assume  $|\mathcal{B}| = 2$ . Without loss of generality,  $\mathcal{B} = \{e_1e_3e_4e_6, e_2e_3e_4e_5\}$ .

1229 Then  $A$  is projectively equivalent to

$$1230 \quad A' = \begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & b \\ 0 & 0 & 0 & 1 & 1 & c \end{pmatrix} \end{matrix}$$

1231 where  $b$  and  $c$  are nonzero, distinct, and not equal to 1. Let

$$1232 \quad T = \begin{pmatrix} -b & -1 & 1 & 0 \\ c & 1 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

1233 The determinant of  $T$  is  $b - c$ , so  $T$  is nonsingular. Now

$$1234 \quad TA' = \begin{pmatrix} -b & -1 & 1 & 0 & 0 & 0 \\ c & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & -1 & -b \\ 0 & 0 & 0 & 1 & 1 & c \end{pmatrix}$$

1235 which has the desired canonical form after column scaling.  $\square$

1236 **Lemma 4.18.** *Let  $(2C_4'', \mathcal{B})$  be a biased graph with no balanced 2-cycle and let*  
 1237  *$A$  be an  $\mathbb{F}$ -matrix representing  $L(2C_4'', \mathcal{B})$ . Then  $A$  is projectively equivalent to*  
 1238 *a canonical lift matrix particular to  $(2C_4'', \mathcal{B})$ .*

1239 *Proof.* We may assume the edges of  $2C_4''$  are labeled as in Figure 14. If  $|\mathcal{B}| = 2$ ,  
 1240 then there is a  $\text{GF}(2)^+$ -gain function  $\gamma$  realizing  $(2C_4'', \mathcal{B})$ . Thus  $L(2C_4'', \mathcal{B})$  is  
 1241 binary, represented by  $A_L(2C_4'', \gamma)$ , so  $L(2C_4'', \mathcal{B})$  has a projectively unique  
 1242 representation over every field, and the result follows. So now assume that

1243 either  $\mathcal{B} = \emptyset$  or, without loss of generality,  $\mathcal{B} = \{e_1 e_3 e_4 e_6\}$ . Since  $\{e_1, e_2, e_5, e_6\}$   
 1244 is a circuit,  $A$  is projectively equivalent to the matrix

$$1245 \quad A' = \begin{pmatrix} e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & a \\ 0 & 0 & 1 & 0 & 1 & b \\ 0 & 0 & 0 & 1 & 1 & b \end{pmatrix}$$

1246 for some  $a, b \in \mathbb{F}$ , where  $a$  and  $b$  are distinct, neither  $a$  nor  $b$  is 1,  $b \neq 0$ , and  
 1247  $a = 0$  if and only if  $|\mathcal{B}| = 1$ . Let

$$1248 \quad T = \begin{pmatrix} 0 & 1-b & 0 & 0 \\ b-a & 1-b & a-1 & 0 \\ a-b & b-1 & 0 & 1-a \\ 0 & 0 & 1-a & 0 \end{pmatrix}.$$

1249 Then  $\det(T) = (a-1)^2(a-b)(b-1) \neq 0$ , so  $T$  is nonsingular, and


$$1250 \quad TA' = \begin{pmatrix} 0 & 1-b & 0 & 0 & 1-b & a-ab \\ b-a & 1-b & a-1 & 0 & 0 & 0 \\ a-b & b-1 & 0 & 1-a & 0 & 0 \\ 0 & 0 & 1-a & 0 & 1-a & b-ab \end{pmatrix}.$$

1251 After scaling columns appropriately (and appending a fifth row obtained by  
 1252 negating the sum of rows 2, 3, and 4, if desired) this is a canonical lift matrix  
 1253 particular to  $(2C_4'', \mathcal{B})$ .  $\square$

1254 For the almost-balanced case of Theorem 2 we need the result analogous  
 1255 to the previous lemmas for one more biased graph. Recall that the graph  
 1256 obtained from  $2C_3$  by deleting an edge is denoted  $2C_3 \setminus e$ .

1257 **Lemma 4.19.** *Let  $(2C_3 \setminus e, \mathcal{B})$  be a biased graph with no balanced 2-cycle and*  
 1258 *let  $A$  be an  $\mathbb{F}$ -matrix representing  $M(2C_3 \setminus e, \mathcal{B})$ . Then  $A$  is projectively equiv-*  
 1259 *alent to a canonical lift matrix particular to  $(2C_3 \setminus e, \mathcal{B})$  and  $A$  is projectively*  
 1260 *equivalent to a canonical frame matrix particular to  $(2C_3 \setminus e, \mathcal{B})$  or to a roll-up*  
 1261 *of  $(2C_3 \setminus e, \mathcal{B})$ .*

1262 *Proof.* Assume  $2C_3 \setminus e$  is labeled as in Figure 15. Then  $\{e_1, e_2, e_3\}$  is a basis,  
 1263 so we may assume the first three columns of  $A$  are labelled  $e_1, e_2, e_3$ , and that

Figure 15:  $2C_3 \setminus e$ .

1264 these columns form an identity matrix. Hence  $A$  is projectively equivalent to  
 1265 the matrix

$$1266 \quad A' = \begin{pmatrix} & e_1 & e_2 & e_3 & e_4 & e_5 \\ 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & a \\ 0 & 0 & 1 & 1 & b \end{pmatrix}$$

1267 for some  $a, b \in \mathbb{F}$ . Since  $\{e_1, e_2, e_5\}$  is not a circuit,  $b \neq 0$ , and since  $\{e_3, e_4, e_5\}$   
 1268 is not a circuit,  $a \neq 1$ . Choose an element  $t \neq -1$  and let

$$1269 \quad T = \begin{pmatrix} t & 1 & -(a+t)/b \\ 1 & -1 & 0 \\ 0 & 0 & (a-1)/b \end{pmatrix}.$$

1270 The determinant of  $T$  is  $(1-a)(t+1)/b$ , so  $T$  is nonsingular, and

$$1271 \quad TA' = \begin{pmatrix} t & 1 & -(a+t)/b & (b-a+t(b-1))/b & 0 \\ 1 & -1 & 0 & 0 & 1-a \\ 0 & 0 & (a-1)/b & (a-1)/b & a-1 \end{pmatrix}.$$

1272 After scaling columns so that every entry in rows 2 and 3 is  $\pm 1$ , this is a  
 1273 canonical lift matrix particular to  $(2C_3 \setminus e, \mathcal{B})$ , where the first row is the “gains  
 1274 row” indexed by  $v_0$  and rows 2 and 3 are indexed by vertices  $v_2$  and  $v_3$ , respec-  
 1275 tively. Now re-index the first row as  $v_1$ . Taking  $t = 0$  yields a canonical frame  
 1276 matrix particular to a roll-up of  $(2C_3 \setminus e, \mathcal{B})$ . If  $\mathbb{F} - \{0, -1, -a, (b-a)/(1-b)\}$   
 1277 is nonempty, then choosing an element from this set for  $t$  yields a canonical  
 1278 frame matrix particular to  $(2C_3 \setminus e, \mathcal{B})$ .  $\square$

## 1279 5 Main results

### 1280 5.1 Projectively equivalent canonical representations arise 1281 from switching equivalent gain graphs

1282 A biased graph representing a 3-connected matroid is 2-connected and has  
1283 no 2-separation with one side inducing a balanced biased subgraph. Thus  
1284 Theorem 1 follows immediately from Theorems 5.1 and 5.4 below.

1285 **Theorem 5.1.** *Let  $(G, \mathcal{B})$  be a loopless, 2-connected, properly unbalanced bi-*  
1286 *ased graph. Let  $\mathbb{F}$  be a field.*

- 1287 1. *The canonical frame matrices given by two  $\mathbb{F}^\times$ -gain functions  $\varphi$  and  $\psi$*   
1288 *realizing  $(G, \mathcal{B})$  are projectively equivalent if and only if  $\varphi$  and  $\psi$  are*  
1289 *switching equivalent.*
- 1290 2. *The canonical lift matrices given by two  $\mathbb{F}^+$ -gain functions  $\varphi$  and  $\psi$*   
1291 *realizing  $(G, \mathcal{B})$  are projectively equivalent if and only if  $\varphi$  and  $\psi$  are*  
1292 *switching-and-scaling equivalent.*
- 1293 3. *Let  $\varphi$  be an  $\mathbb{F}^\times$ -realization of  $(G, \mathcal{B})$  and let  $\psi$  be an  $\mathbb{F}^+$ -realization of*  
1294  *$(G, \mathcal{B})$ . Then  $A_F(G, \varphi)$  and  $A_L(G, \psi)$  are not projectively equivalent.*

1295 *Proof.* For part (1) (respectively, part (2)), if  $\varphi$  and  $\psi$  are switching (resp.  
1296 switching-and-scaling) equivalent then  $A_F(G, \varphi)$  and  $A_F(G, \psi)$  (resp.  $A_L(G, \varphi)$   
1297 and  $A_L(G, \psi)$ ) are projectively equivalent by Proposition 4.2.

1298 For the converse of part (1) (resp. part (2)) assume that  $\varphi$  and  $\psi$  are  
1299  $\mathbb{F}^\times$ -realizations (resp.  $\mathbb{F}^+$ -realizations) that are not switching (resp. switching-  
1300 and-scaling) equivalent. By Theorem 6, there is a minor  $(H, \mathcal{S})$  of  $(G, \mathcal{B})$   
1301 such that either  $(H, \mathcal{S}) \in \mathcal{G}_0$  with  $\varphi|_H$  and  $\psi|_H$  switching inequivalent or  
1302  $(H, \mathcal{S}) \cong \mathbf{U}_2$  (resp.  $(H, \mathcal{S}) \cong \mathbf{U}_3$ ) with  $\varphi|_H$  and  $\psi|_H$  switching (resp. switching-  
1303 and-scaling) inequivalent on the 2-cycle of  $\mathbf{U}_2$  (resp. on the theta subgraph of  
1304  $\mathbf{U}_3$ ). By Lemmas 4.4, 4.7, 4.10, and 4.12,  $A_F(H, \varphi|_H)$  and  $A_F(H, \psi|_H)$  are not  
1305 projectively equivalent (resp. by Lemmas 4.5, 4.8, 4.11, and 4.13,  $A_L(H, \varphi|_H)$   
1306 and  $A_L(H, \psi|_H)$  are not projectively equivalent). Thus  $A_F(G, \varphi)$  and  $A_F(G, \psi)$   
1307 (resp.  $A_L(G, \varphi)$  and  $A_L(G, \psi)$ ) are not projectively equivalent.

1308 For part (3), first observe that if  $(G, \mathcal{B})$  is not tangled, then  $F(G, \mathcal{B}) \neq$   
1309  $L(G, \mathcal{B})$  so  $A_F(G, \varphi)$  and  $A_L(G, \psi)$  are certainly not projectively equivalent. So  
1310 assume  $(G, \mathcal{B})$  is tangled. Then by Corollary 3.6  $(G, \mathcal{B})$  has a minor  $(H, \mathcal{S}) \in \mathcal{G}_0$   
1311 where  $H$  is either  $2C_3$  or  $K_4$ . Thus by Lemmas 4.6 and 4.9  $A_F(G, \varphi)$  and  
1312  $A_L(G, \psi)$  are not projectively equivalent.  $\square$

1313 **Derived gain functions.** Recall that when  $(G, \mathcal{B})$  has no two vertex-disjoint  
 1314 unbalanced cycles,  $F(G, \mathcal{B}) = L(G, \mathcal{B})$ , and we denote this matroid  $M(G, \mathcal{B})$ .  
 1315 Recall also that for an  $\mathbb{F}^+$ -gain function  $\varphi$ , in the case that  $(G, \mathcal{B}_\varphi)$  has a  
 1316 balancing vertex  $u$  we assume that  $A_L(G, \varphi)$  is of full rank such that removing  
 1317 its row corresponding to the gains assigned by  $\varphi$  leaves the oriented incidence  
 1318 matrix of  $G$  minus row  $u$  (as described in Section 2.6). Recall also that for every  
 1319 almost-balanced biased graph  $(G, \mathcal{B})$ , there is a family of biased graphs  $\mathcal{R}_{(G, \mathcal{B})}$ ,  
 1320 each member of which represents  $M(G, \mathcal{B})$  as a frame matroid, and there is  
 1321 a uniquely chosen member  $(\widehat{G}, \widehat{\mathcal{B}})$  of  $\mathcal{R}_{(G, \mathcal{B})}$  such that all other members of  
 1322  $\mathcal{R}_{(G, \mathcal{B})}$  are obtained from  $(\widehat{G}, \widehat{\mathcal{B}})$  as roll-ups (Section 2.5).

1323 Let  $(G, \mathcal{B})$  be a biased graph with a joint. Then no  $\text{GF}(2)^\times$ -gain function  
 1324 can realize  $(G, \mathcal{B})$ , for the trivial reason that  $\text{GF}(2)^\times$  has no non-identity ele-  
 1325 ment. Aside from those over  $\text{GF}(2)$ , there is a very close relationship between  
 1326 gain functions from the additive and multiplicative groups of a field realiz-  
 1327 ing almost-balanced biased graphs. Let  $(G, \mathcal{B})$  be an almost-balanced biased  
 1328 graph, with balancing vertex  $u$  after deleting its set  $J$  of joints. Every gain  
 1329 function realizing  $(G, \mathcal{B})$  is switching equivalent to a gain function assigning  
 1330 the group identity element to each link not incident  $u$ , obtained by normalizing  
 1331 on a spanning tree of  $G - u$ . So let  $T$  be a spanning tree of  $G - u$ , and let  $\varphi$  be  
 1332  $T$ -normalized  $\mathbb{F}^\times$ -gain function realizing  $(G, \mathcal{B})$ . Then there is a  $T$ -normalized  
 1333  $\mathbb{F}^+$ -gain function  $\varphi^+$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  obtained from  $\varphi$  up to loops by simply  
 1334 replacing the multiplicative identity with the additive identity. That is, set  
 1335  $\varphi^+(e) = 0$  if  $e$  is a link not incident to  $u$  and  $\varphi^+(e) = \varphi(e)$  if  $e$  is a link incident  
 1336 to  $u$ . To complete the definition of  $\varphi^+$ , simply set  $\varphi^+(e) = 0$  if  $e$  is joint not  
 1337 incident to  $u$  or a balanced loop, and  $\varphi^+(e) = -1$  if  $e$  is a joint incident to  $u$ .  
 1338 Call  $\varphi^+$  the  $\mathbb{F}^+$ -gain function *derived from*  $\varphi$ .

1339 For each unbalancing class  $U$  of  $\Sigma(u)$ , let us denote by  $(G_U, \mathcal{B}_U)$  the roll-up  
 1340 of  $(\widehat{G}, \widehat{\mathcal{B}})$  in which  $U$  is a set of joints. Let  $T_U$  be a spanning tree of  $\widehat{G}$  containing  
 1341 an edge in  $U$ , and let  $T$  be the spanning tree of  $G - u$  obtained from  $T_U$  by  
 1342 deleting its edge incident to  $u$ . Let  $\psi$  be a  $T_U$ -normalized  $\mathbb{F}^+$ -gain function  
 1343 realizing  $(\widehat{G}, \widehat{\mathcal{B}})$ . Assume  $\mathbb{F}$  is not  $\text{GF}(2)$ , and choose an element  $a \in \mathbb{F}^\times$  that  
 1344 is not 1. Then there is a  $T$ -normalized  $\mathbb{F}^\times$ -gain function  $\psi^\times$  realizing  $(G_U, \mathcal{B}_U)$   
 1345 obtained from  $\psi$  up to loops by simply replacing the additive identity with  
 1346 the multiplicative identity. That is, set  $\psi^\times(e) = 1$  if  $e$  is a link not incident to  
 1347  $u$  and  $\psi^\times(e) = \psi(e)$  if  $e$  is a link incident to  $u$ . Every link  $e$  incident to  $u$  in  
 1348  $G_U$  satisfies  $\psi(e) \neq 0$ , and so satisfies  $\psi(e) \in \mathbb{F}^\times$ . Complete the definition of  
 1349  $\psi^\times$  by simply setting  $\psi^\times(e) = 1$  if  $e$  is a balanced loop and  $\psi^\times(e) = a$  if  $e$  is a  
 1350 joint. Call  $\psi^\times$  the  $\mathbb{F}^\times$ -gain function *derived from*  $\psi$ .

1351 **Lemma 5.2.** *Let  $(G, \mathcal{B})$  be an almost-balanced biased graph and let  $\mathbb{F}$  be a*

1352 field other than  $\text{GF}(2)$ . Let  $J$  be the set of joints of  $(G, \mathcal{B})$ . Assume  $(G, \mathcal{B}) \setminus J$   
 1353 has a unique balancing vertex  $u$ , and let  $T$  be a spanning tree of  $G - u$ .

1354 1. Let  $\varphi$  be a  $T$ -normalized  $\mathbb{F}^\times$ -gain function realizing a roll-up  $(G_U, \mathcal{B}_U)$  of  
 1355  $(\widehat{G}, \widehat{\mathcal{B}})$ . Then  $A_L(\widehat{G}, \varphi^+)$  is obtained from  $A_F(G_U, \varphi)$  by scaling columns.

1356 2. Let  $\psi$  be a  $T$ -normalized  $\mathbb{F}^+$ -gain function realizing  $(\widehat{G}, \widehat{\mathcal{B}})$ . Let  $U$  be the  
 1357 (possibly empty) unbalancing class of  $\Sigma(u)$  for which  $\psi(U) = \{0\}$ . Then  
 1358  $A_F(G_U, \psi^\times)$  is obtained from  $A_L(\widehat{G}, \psi)$  by scaling columns.

1359 *Proof.* (1) Since  $\varphi$  is  $T$ -normalized,  $\varphi(e) = 1$  for each link  $e$  that is not incident  
 1360 to  $u$ . We may assume that all edges incident to  $u$  are directed into  $u$ . Let  $A$  be  
 1361 the matrix obtained from  $A_F(G_U, \varphi)$  by re-indexing row  $u$  as  $v_0$  (the ‘‘gains’’  
 1362 row) and scaling columns so that each column that is nonzero in just one row  
 1363 has that nonzero entry equal to  $-1$ , and columns with a nonzero entry in row  
 1364  $v_0$  have either all other entries 0 or have other nonzero entry equal to  $-1$ .  
 1365 Because  $\varphi$  assigns 1 to every link not incident to  $u$ ,  $A$  is a canonical lift matrix  
 1366 particular to  $(\widehat{G}, \widehat{\mathcal{B}})$ , where each edge in  $\Sigma(u)$  is a link directed out from  $u$ .  
 1367 Moreover, for each edge  $e \in E(G)$ , the entry in row  $v_0$ , column  $e$  of  $A$  is equal  
 1368 to  $\varphi^+(e)$ , so  $A = A_L(\widehat{G}, \varphi^+)$ .

1369 (2) Since  $\psi$  is  $T$ -normalized,  $\psi(e) = 0$  for each edge  $e$  that is not incident  
 1370 to  $u$ . We may assume that all links incident to  $u$  in  $G$  are directed out from  $u$ .  
 1371 Since by assumption the rows of  $A_L(\widehat{G}, \psi)$  are indexed by  $v_0$  (the ‘‘gains row’’)  
 1372 and  $V(\widehat{G}) - u$ , every column of  $A_L(\widehat{G}, \psi)$  has at most two nonzero entries.  
 1373 Let  $A$  be the matrix obtained from  $A_L(\widehat{G}, \psi)$  by re-indexing row  $v_0$  as row  $u$   
 1374 and scaling columns so that each column that has exactly two nonzero entries  
 1375 with a nonzero entry in row  $u$  has its other nonzero entry equal to 1, and  
 1376 every column with just one nonzero entry has its nonzero entry equal to  $1 - a$ ,  
 1377 where  $a$  is the chosen element of  $\mathbb{F}^\times$  different than 1 that  $\psi^\times$  assigns to joints.  
 1378 Since  $(\widehat{G}, \widehat{\mathcal{B}})$  has  $u$  as a balancing vertex, each column  $e$  of  $A$  with 0 in row  $u$   
 1379 has either exactly two nonzero entries, which are 1 and  $-1$ , and these appear  
 1380 in rows  $v, w$  when  $e$  has endpoints  $v, w$  and neither  $v$  nor  $w$  is equal to  $u$ , or  
 1381 column  $e$  at most one nonzero entry, which is  $1 - a$ , appearing in row  $v$  when  
 1382  $e \in U$  with endpoints  $u, v$  in  $\widehat{G}$ . Thus  $A$  is a canonical frame matrix particular  
 1383 to  $(G_U, \mathcal{B}_U)$  where each edge in  $\Sigma(u) - U$  is a link directed into  $u$  and each  
 1384 element in  $U$  is a joint. Moreover, by definition the gain function  $\psi^\times$  realizes  
 1385  $(G_U, \mathcal{B}_U)$  and  $A = A_F(G_U, \psi^\times)$ .  $\square$

1386 Let  $(G, \mathcal{B})$  be an almost-balanced biased graph with a unique balancing  
 1387 vertex  $u$  after deleting its joints. Since we may always assume that all links  
 1388 incident to  $u$  are directed either into or out from  $u$ , and every gain function

1389 realizing  $(G, \mathcal{B})$  is switching equivalent to a gain function assigning the identity  
 1390 element to each link not incident to  $u$ , we may define a *derived* gain function  
 1391 from any  $\mathbb{F}^\times$ - or  $\mathbb{F}^+$ -gain function, by first switching appropriately. Further, for  
 1392 each  $\mathbb{F}^+$ -gain function realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  and each unbalancing class  $U \in \Sigma(u)$ ,  
 1393 we may always switch to obtain an  $\mathbb{F}^+$ -gain function  $\varphi$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  with the  
 1394 property that for each edge  $e \in U$ ,  $\varphi(e) = 0$ . Thus, with this extension of the  
 1395 notion of a derived gain function, Proposition 4.2 and Lemma 5.2 immediately  
 1396 yield the following.

1397 **Corollary 5.3.** *Let  $(G, \mathcal{B})$  be an almost-balanced biased graph with a unique*  
 1398 *balancing vertex  $u$  after deleting its set of joints, and let  $\mathbb{F}$  be a field other than*  
 1399  *$\text{GF}(2)$ .*

- 1400 1. *For every  $\mathbb{F}^+$ -gain function  $\varphi$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$ , and every unbalancing*  
 1401 *class  $U \in \Sigma(u)$ , there is a derived  $\mathbb{F}^\times$ -gain function  $\varphi^\times$  realizing the*  
 1402 *roll-up  $(G_U, \mathcal{B}_U)$  of  $(\widehat{G}, \widehat{\mathcal{B}})$  such that  $A_L(\widehat{G}, \varphi)$  and  $A_F(G_U, \varphi^\times)$  are pro-*  
 1403 *jectively equivalent.*
- 1404 2. *For every  $\mathbb{F}^\times$ -gain function  $\psi$  realizing a roll-up  $(G_U, \mathcal{B}_U)$  of  $(\widehat{G}, \widehat{\mathcal{B}})$ , there*  
 1405 *is a derived  $\mathbb{F}^+$ -gain function  $\psi^+$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  for which  $A_F(G_U, \psi)$*   
 1406 *and  $A_L(\widehat{G}, \psi^+)$  are projectively equivalent.*
- 1407 3. *There is an  $\mathbb{F}^\times$ -gain function realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  if and only if there is an*  
 1408  *$\mathbb{F}^+$ -gain function  $\varphi$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  for which  $\varphi(e) \neq 0$  for each edge in*  
 1409  *$\Sigma(u)$ .*

1410 Equipped with the above tool, we can now state and prove a result analo-  
 1411 gous to Theorem 5.1 for almost-balanced biased graphs.

1412 **Theorem 5.4.** *Let  $(G, \mathcal{B})$  be a 2-connected, almost-balanced biased graph with*  
 1413 *a unique balancing vertex  $u$  after deleting its joints, and with no vertical 2-*  
 1414 *separation with one side balanced. Let  $\mathbb{F}$  be a field.*

- 1415 1. *The canonical lift matrices given by two  $\mathbb{F}^+$ -gain functions  $\varphi$  and  $\psi$*   
 1416 *realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  are projectively equivalent if and only if  $\varphi$  and  $\psi$  are*  
 1417 *switching-and-scaling equivalent.*
- 1418 2. *Let  $U$  and  $W$  be unbalancing classes of  $\Sigma(u)$ . Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^\times$ -gain*  
 1419 *functions realizing  $(G_U, \mathcal{B}_U)$  and  $(G_W, \mathcal{B}_W)$ , respectively. The canoni-*  
 1420 *cal frame matrices given by  $\varphi$  and  $\psi$  are projectively equivalent if and*  
 1421 *only if their derived gain functions  $\varphi^+$  and  $\psi^+$  are switching-and-scaling*  
 1422 *equivalent.*

- 1423 3. Let  $U$  be an unbalancing class of  $\Sigma(u)$ . The canonical lift matrix given  
 1424 by an  $\mathbb{F}^+$ -gain function  $\varphi$  realizing  $(\widehat{G}, \widehat{\mathcal{B}})$  and the canonical frame ma-  
 1425 trix given by an  $\mathbb{F}^\times$ -gain function  $\psi$  realizing  $(G_U, \mathcal{B}_U)$  are projectively  
 1426 equivalent if and only if  $\varphi$  and the derived gain function  $\psi^+$  are switching-  
 1427 and-scaling equivalent.
- 1428 4. Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^+$ - and  $\mathbb{F}^\times$ -gain functions, respectively, realizing  $(\widehat{G}, \widehat{\mathcal{B}})$ .  
 1429 The canonical lift matrix given by  $\varphi$  and the canonical frame matrix  
 1430 given by  $\psi$  are projectively equivalent if and only if  $\varphi$  and the derived  
 1431 gain function  $\psi^+$  are switching-and-scaling equivalent.

1432 *Proof.* (1) Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^+$ -gain functions realizing  $(\widehat{G}, \widehat{\mathcal{B}})$ . If  $\varphi$  and  $\psi$   
 1433 are switching-and-scaling equivalent then by Proposition 4.2  $A_L(\widehat{G}, \varphi)$  and  
 1434  $A_L(\widehat{G}, \psi)$  are projectively equivalent.

1435 Conversely, suppose for a contradiction that  $A_L(\widehat{G}, \varphi)$  and  $A_L(\widehat{G}, \psi)$  are  
 1436 projectively equivalent but that  $\varphi$  and  $\psi$  are not switching-and-scaling equiv-  
 1437 alent. We may assume that all edges incident to  $u$  have  $u$  as their tail. By  
 1438 switching we may assume that  $\varphi(e) = \psi(e) = 0$  for each edge  $e$  not incident to  
 1439  $u$ . We may further assume that  $A_L(\widehat{G}, \varphi)$  and  $A_L(\widehat{G}, \psi)$  are both of full rank  
 1440 with rows indexed by  $v_0 \cup (V(G) - u)$ , where  $v_0 \notin V(G)$  is the ‘‘gains row’’.  
 1441 Put  $A = A_L(\widehat{G}, \varphi)$  and  $B = A_L(\widehat{G}, \psi)$ , and suppose  $T$  is a nonsingular matrix  
 1442 such that  $TAS = B$ , where  $S$  is a nonsingular diagonal column-scaling matrix.  
 1443 Let the rows and columns of  $T$  be indexed by  $v_0 \cup (V(G) - u)$  according to  
 1444 the rows of  $A$ . By Lemma 4.3 either  $T$  has an entry on its main diagonal that  
 1445 is not 1 or  $T$  has a nonzero entry off its main diagonal, in either case in a row  
 1446 other than  $v_0$ . Suppose first  $T$  has all entries off its main diagonal equal to  
 1447 0, aside from those in row  $v_0$ . Let  $U$  be the  $|V(G)| \times |V(G)|$  diagonal matrix  
 1448 with rows and columns indexed by  $v_0 \cup V(G) - u$  in which entry  $U_{v_0 v_0} = 1$   
 1449 and entry  $U_{vv} = a^{-1}$  if entry  $T_{vv} = a$ . Since  $T$  is non-singular no such entry  
 1450 is 0. Removing the row and column of  $UT$  indexed by  $v_0$  leaves an identity  
 1451 matrix and  $(UT)AR = B$ , where  $R$  is an appropriate diagonal matrix scaling  
 1452 the columns of  $(UT)A$ . Thus  $A$  and  $B$  are projectively equivalent, contrary to  
 1453 assumption. So assume there is a nonzero element off the main diagonal of  $T$   
 1454 in a row other than  $v_0$ . Suppose the entry in row  $x$ , column  $y$  is nonzero, where  
 1455  $x \neq y$  and  $x \neq v_0$ . Since  $G$  is 2-connected and has no vertical 2-separation with  
 1456 one side balanced, there is an unbalanced cycle  $C$  avoiding  $x$  while containing  
 1457  $y$ . Let  $f, f'$  be the edges of  $C$  incident to  $u$ . Then  $\varphi(f) \neq \varphi(f')$ . Denote by  
 1458  $T_x$  row  $x$  of  $T$  and by  $A_e$  column  $e$  of  $A$ . Consider the equations  $T_x \cdot A_e = B_{xe}$   
 1459 given by each of the dot products of row  $x$  of  $T$  with column  $e$  of  $A$ , for each  
 1460  $e \in E(C)$ . Since  $C$  avoids  $x$ , for each edge  $e \in E(C)$  entry  $B_{xe}$  is zero. There



1461 are precisely two nonzero entries in each column  $e$  of  $A$  with  $e \in E(C)$ , and  
 1462 other than columns  $f$  and  $f'$  one of these two entries is 1 and the other is  $-1$ .  
 1463 Thus the system of equations  $\{T_x \cdot A_e = 0 : e \in E(C)\}$  imply  $\varphi(f) = \varphi(f')$ , a  
 1464 contradiction.

1465 (2) If  $\varphi^+$  and  $\psi^+$  are switching-and-scaling equivalent, then by Proposition  
 1466 4.2  $A_L(\widehat{G}, \varphi^+)$  and  $A_L(\widehat{G}, \psi^+)$  are projectively equivalent. Hence by Corollary  
 1467 5.3  $A_F(G_U, \varphi)$  and  $A_F(G_W, \psi)$  are projectively equivalent. Conversely, sup-  
 1468 pose  $A_F(G_U, \varphi)$  and  $A_F(G_W, \psi)$  are projectively equivalent. Then by Corollary  
 1469 5.3  $A_L(\widehat{G}, \varphi^+)$  and  $A_L(\widehat{G}, \psi^+)$  are projectively equivalent, and so by statement  
 1470 (1)  $\varphi^+$  and  $\psi^+$  are switching-and-scaling equivalent.

1471 The proofs of statements (3) and (4) are straightforward modifications of  
 1472 the proof of (2).  $\square$

## 1473 5.2 Matrix representations arise from biased graph rep- 1474 resentations

1475 Theorem 2 is an immediate consequence of Theorem 5.5 below.

1476 **Theorem 5.5.** *Let  $M$  be a 3-connected matroid of rank greater than two, and*  
 1477 *let  $\mathbb{F}$  be a field. Let  $A$  be a matrix over  $\mathbb{F}$  representing  $M$  and let  $(G, \mathcal{B})$  be a*  
 1478 *biased graph representing  $M$ .*

1479 1. *If  $(G, \mathcal{B})$  is properly unbalanced then exactly one of the following holds.*

1480 (i)  *$A$  is projectively equivalent to a canonical lift matrix particular to*  
 1481  *$(G, \mathcal{B})$ , or*

1482 (ii)  *$A$  is projectively equivalent to a canonical frame matrix particular*  
 1483 *to  $(G, \mathcal{B})$ .*

1484 2. *If  $(G, \mathcal{B})$  is almost-balanced then each of the following hold, unless  $\mathbb{F}$  is*  
 1485  *$\text{GF}(2)$ .*

1486 (i)  *$A$  is projectively equivalent to a canonical lift matrix particular to*  
 1487  *$(\widehat{G}, \widehat{\mathcal{B}})$ , and*

1488 (ii)  *$A$  is projectively equivalent to a canonical frame matrix particular*  
 1489 *to each roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$ .*

1490 *In the case  $\mathbb{F}$  is  $\text{GF}(2)$ , (i) holds.*

1491 When  $(G, \mathcal{B})$  is almost-balanced, either  $(G, \mathcal{B}) = (\widehat{G}, \widehat{\mathcal{B}})$  or  $(G, \mathcal{B})$  is a  
 1492 roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$ . Thus Theorem 5.5 says that in the almost-balanced case,

1493  $A$  is projectively equivalent to a canonical lift matrix particular to  $(G, \mathcal{B})$  if  
 1494  $(G, \mathcal{B}) = (\widehat{G}, \widehat{\mathcal{B}})$ , while  $A$  is projectively equivalent to a canonical frame matrix  
 1495 particular to  $(G, \mathcal{B})$  if  $(G, \mathcal{B})$  is a roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$ .

1496 Theorem 5.5 follows immediately from Theorems 5.6 and 5.9 below.

1497 **Theorem 5.6.** *Let  $M$  be a matroid represented by a 2-connected, properly*  
 1498 *unbalanced biased graph  $(G, \mathcal{B})$ . Let  $\mathbb{F}$  be a field and let  $A$  be a matrix over  $\mathbb{F}$*   
 1499 *representing  $M$ . Exactly one of the following holds:*

1500 (i)  *$A$  is projectively equivalent to a canonical lift matrix particular to  $(G, \mathcal{B})$ ,*  
 1501 *or*

1502 (ii)  *$A$  is projectively equivalent to a canonical frame matrix particular to*  
 1503  *$(G, \mathcal{B})$ .*

1504 We will require the following fact on several occasions. It follows immedi-  
 1505 ately from the fact that a pair of edges incident to a vertex of degree two form  
 1506 a series pair in the matroid.

1507 **Lemma 5.7.** *Let  $(H, \mathcal{S})$  be a subdivision of  $(G, \mathcal{B})$ . Let  $\mathbb{F}$  be a field and let*  
 1508  *$\Gamma \in \{\mathbb{F}^+, \mathbb{F}^\times\}$ .*

1509 1. *The  $\mathbb{F}$ -matrix representations of  $F(H, \mathcal{S})$  are in one-to-one correspon-*  
 1510 *dence with the  $\mathbb{F}$ -matrix representations of  $F(G, \mathcal{B})$  up to projective equiv-*  
 1511 *alence.*

1512 2. *The  $\mathbb{F}$ -matrix representations of  $L(H, \mathcal{S})$  are in one-to-one correspon-*  
 1513 *dence with the  $\mathbb{F}$ -matrix representations of  $L(G, \mathcal{B})$  up to projective equiv-*  
 1514 *alence.*

1515 3. *The  $\Gamma$ -realizations of  $(H, \mathcal{S})$  are in one-to-one correspondence with the*  
 1516  *$\Gamma$ -realizations of  $(G, \mathcal{B})$  up to switching (resp. switching-and-scaling).*

1517 We also need the following more technical fact to prove Theorem 5.6.

1518 **Lemma 5.8.** *Let  $(G, \mathcal{B})$  be a connected biased graph with a joint  $e$  such that*  
 1519  *$(G, \mathcal{B}) \setminus e$  is a biased  $2C_3$  with no balanced 2-cycle or a biased  $K_4$  with no*  
 1520 *balanced triangle. Let  $\mathbb{F}$  be a field, and let  $\varphi$  be an  $\mathbb{F}^\times$ - or  $\mathbb{F}^+$ -gain function on*  
 1521  *$G$ .*

1522 1. *If  $A_F(G \setminus e, \varphi)$  represents  $M((G, \mathcal{B}) \setminus e)$  then  $A_F(G \setminus \ell, \varphi)$  does not extend*  
 1523 *to an  $\mathbb{F}$ -representation of  $L(G, \mathcal{B})$ .*

1524 2. If  $A_L(G \setminus e, \varphi)$  represents  $M((G, \mathcal{B}) \setminus e)$  then  $A_L(G \setminus \ell, \varphi)$  does not extend  
 1525 to an  $\mathbb{F}$ -representation of  $F(G, \mathcal{B})$ .

1526 *Proof.* We give a detailed proof for the case in which  $(G, \mathcal{B}) \setminus e$  is a biased  
 1527  $2C_3$ . The case for which  $(G, \mathcal{B}) \setminus e$  is a biased  $K_4$  follows from  $\Delta$ - $Y$  and  $Y$ - $\Delta$   
 1528 exchanges, by Propositions 2.8 and 2.11.

1529 (1) Suppose for a contradiction that there is a matrix  $A$  over  $\mathbb{F}$  representing  
 1530  $L(G, \mathcal{B})$  such that removing column  $e$  from  $A$  yields the matrix  $A_F(G \setminus e, \varphi)$ .  
 1531 We may assume that  $A$  has full rank, and so has three rows. Since  $e$  is  
 1532 not a loop of  $L(G, \mathcal{B})$ , column  $e$  of  $A$  is nonzero. If column  $e$  has just one  
 1533 nonzero entry, then  $A$  is an  $\mathbb{F}$ -representation of a matroid  $F(\Omega)$  where  $\Omega$  is  
 1534 obtained from  $(G, \mathcal{B}) \setminus e$  by adding a joint to a vertex. But comparing circuits  
 1535 we see that  $F(\Omega) \neq L(G, \mathcal{B})$ , a contradiction. If column  $e$  has exactly two  
 1536 nonzero entries, then  $A$  is an  $\mathbb{F}$ -representation of a matroid  $F(\Omega)$ , where  $\Omega$   
 1537 is obtained from  $(G, \mathcal{B}) \setminus e$  by adding a link. But again comparing circuits we  
 1538 see that  $F(\Omega) \neq L(G, \mathcal{B})$ , a contradiction. So finally suppose column  $e$  has  
 1539 three nonzero entries. Let  $X$  be an unbalanced 2-cycle of  $\Omega$ . Then  $X \cup e$  is  
 1540 a circuit of  $L(G, \mathcal{B})$  but the columns of  $A$  corresponding to  $X \cup e$  are linearly  
 1541 independent, a contradiction.

1542 (2) Suppose for a contradiction that there is a matrix  $A$  representing  
 1543  $F(G, \mathcal{B})$  such that removing column  $e$  from  $A$  yields  $A_L(G \setminus e, \varphi)$ . Let  $V(G) =$   
 1544  $\{v_1, v_2, v_3\}$  where  $e$  is incident to  $v_1$ . Then  $A \setminus e = A_L(G \setminus e, \varphi)$  has rows in-  
 1545 dexed by  $v_0 \cup V(G)$  where  $v_0 \notin V(G)$  is the “gains row” and removing row  $v_0$   
 1546 from  $A_L(G \setminus e, \varphi)$  leaves the oriented incidence matrix of  $G \setminus e$ . Since  $F(G, \mathcal{B})$   
 1547 has rank three while  $A$  has four rows,  $A$  is not of full rank. Since in  $A \setminus e$   
 1548 row  $v_0$  is not in the span of  $\{v_1, v_2, v_3\}$ , neither is row  $v_0$  in the span of rows  
 1549  $\{v_1, v_2, v_3\}$  in  $A$ . Thus in  $A$  row  $v_3$  is in the span of rows  $v_1$  and  $v_2$ . Since any  
 1550 linear combination of rows  $v_1, v_2$ , and  $v_3$  in  $A$  yields a corresponding linear  
 1551 combination in  $A \setminus e$ , this implies that the entries of  $A$  in rows  $v_1, v_2$ , and  $v_3$  of  
 1552 column  $e$  sum to zero. Put  $A_{v_1e} = a, A_{v_2e} = b$ ; then  $A_{v_3e} = -(a + b)$ .

1553 First suppose that  $a = b = 0$ . As  $e$  is not a loop of  $F(G, \mathcal{B})$ , column  $e$  is  
 1554 nonzero. Thus  $A_{v_0e} \neq 0$ . Let  $X$  be the unbalanced 2-cycle consisting of the  
 1555 pair of edges linking  $v_2$  and  $v_3$ . Then  $X \cup e$  is independent in  $F(G, \mathcal{B})$  but  
 1556 the columns of  $A$  representing  $X \cup e$  are linearly dependent, a contradiction.  
 1557 Next suppose  $a = -b$ . Let  $X$  be the unbalanced 2-cycle consisting of the  
 1558 edges linking  $v_1$  and  $v_3$ . The set  $X \cup e$  is a circuit in  $F(G, \mathcal{B})$  but has columns  
 1559 linearly independent in  $A$ , a contradiction. Finally, suppose none of  $a, b$ , nor  
 1560  $-(a + b)$  are zero. Let  $X$  be the unbalanced 2-cycle consisting of the edges  
 1561 linking  $v_1$  and  $v_2$ . Then  $X \cup e$  is a circuit of  $F(G, \mathcal{B})$  but has columns linearly  
 1562 independent in  $A$ , a contradiction.  $\square$

1563 *Proof of Theorem 5.6.* We show that if  $M = F(G, \mathcal{B})$  then there is an  $\mathbb{F}^\times$ -  
 1564 gain function  $\varphi$  such that  $A$  is projectively equivalent to  $A_F(G, \varphi)$ , that if  
 1565  $M = L(G, \mathcal{B})$  then there is an  $\mathbb{F}^+$ -gain function  $\psi$  such that  $A$  is projectively  
 1566 equivalent to  $A_L(G, \psi)$ , and that if  $(G, \mathcal{B})$  is tangled, so  $F(G, \mathcal{B}) = L(G, \mathcal{B})$ ,  
 1567 then  $A$  is not projectively equivalent to both  $A_F(G, \varphi)$  and  $A_L(G, \psi)$ .

1568 By Theorem 5,  $(G, \mathcal{B})$  contains a subgraph  $\Omega_0$  that is a subdivision of a  
 1569 biased graph in  $\mathcal{T}_0$ . Let  $(G_0, \mathcal{B}_0)$  be the biased subgraph of  $(G, \mathcal{B})$  induced  
 1570 by  $E(\Omega_0)$ , with  $V(G_0) = V(G)$ . Let  $A_0$  be the submatrix of  $A$  consisting of  
 1571 the columns whose elements are in  $E(\Omega_0)$ . By Lemmas 4.14, 4.15, 4.16, 4.17,  
 1572 or 4.18, and Lemma 5.7, there exists an  $\mathbb{F}^\times$ -gain function  $\varphi_0$  such that  $A_0$  is  
 1573 projectively equivalent to  $A_F(G_0, \varphi_0)$ , or there exists an  $\mathbb{F}^+$ -gain function  $\psi_0$   
 1574 such that  $A_0$  is projectively equivalent to  $A_L(G_0, \psi_0)$ , but not both.

1575 Let  $J$  be the set of joints of  $(G, \mathcal{B})$ . Since both  $\Omega_0$  and  $G$  are 2-connected,  
 1576 there is a sequence of 2-connected biased subgraphs  $\Omega_0 \subset \Omega_1 \subset \dots \subset \Omega_n$   
 1577 where  $\Omega_n = (G, \mathcal{B}) \setminus J$  such that for each  $i \in \{0, \dots, n-1\}$  there is a path  $P_i$  in  
 1578  $G$  internally disjoint from  $\Omega_i$  so that  $\Omega_i \cup P_i = \Omega_{i+1}$ . For each  $i \in \{0, \dots, n\}$ ,  
 1579 let  $(G_i, \mathcal{B}_i)$  be the biased subgraph of  $(G, \mathcal{B})$  induced by  $E(\Omega_i)$  with  $V(G_i) =$   
 1580  $V(G)$ . Let  $A_i$  be the submatrix of  $A$  consisting of all rows of  $A$  and precisely  
 1581 those columns representing  $E(\Omega_i)$ . Thus for each  $i$ , in the case  $M = F(G, \mathcal{B})$   
 1582  $A_i$  represents  $F(\Omega_i)$ ; in the case  $M = L(G, \mathcal{B})$ ,  $A_i$  represents  $L(\Omega_i)$ . Induc-  
 1583 tively assume that for some  $i \geq 0$  there exist nonsingular matrices  $T_0, \dots, T_i$ ,  
 1584 nonsingular diagonal matrices  $S_0, \dots, S_i$ , and gain functions  $\varphi_0, \dots, \varphi_i$ , such  
 1585 that either

1586 (1)  $T_j A_j S_j = A_F(G_j, \varphi_j)$  for each  $j \in \{0, \dots, i\}$ , or

1587 (2)  $T_j A_j S_j = A_L(G_j, \psi_j)$  for each  $j \in \{0, \dots, i\}$ .

1588 We will show the same projective equivalence for  $A_{i+1}$ . We first obtain this  
 1589 conclusion in the case that  $P_i$  consists of a single edge. Then if  $P_i$  has length  
 1590 greater than one, the conclusion follows from Lemma 5.7. So suppose  $P_i$   
 1591 consists of a single edge  $e_i$  linking vertices  $u_i, v_i \in V(\Omega_i)$ . We consider cases  
 1592 (1) and (2) above separately.

1593 (1) Consider the matrix  $T_i A_{i+1}$ . Matrix  $A_{i+1}$  has rows indexed by  $V(G)$ ,  
 1594 according to the indexing of the corresponding rows of  $T_i A_i S_i$ , and columns  
 1595 indexed by  $E(G_{i+1})$ . We first show that column  $e_i$  of  $T_i A_{i+1}$  is zero in every  
 1596 row aside from  $u_i$  and  $v_i$ . Suppose for a contradiction that  $e_i$  is nonzero in a  
 1597 row  $x$  of  $T_i A_{i+1}$  differing from  $u_i$  and  $v_i$ . Since  $\Omega_0$  does not have a balancing  
 1598 vertex, neither does  $\Omega_i$ . Thus  $\Omega_i - x$  is unbalanced and connected. Hence there  
 1599 is a subset  $U_i \subseteq E(\Omega_i - x)$  that induces a subgraph of  $\Omega_i$  that is a spanning  
 1600 tree of  $\Omega_i - x$  along with one additional edge whose fundamental cycle with

1601 respect to this tree is unbalanced. Contained in  $U_i \cup e_i$  is a biased subgraph  $C$   
 1602 whose edge set is a circuit of  $F(\Omega_{i+1})$ . The subgraph  $C$  contains  $e_i$  and does  
 1603 not contain  $x$ . Since  $T_i A_i S_i = A_F(G_i, \varphi_i)$ , the columns of  $T_i A_{i+1}$  representing  
 1604  $E(C) - e_i$  are all zero in row  $x$ . Hence while  $C$  is a circuit of  $F(\Omega_{i+1})$  the  
 1605 columns of  $T_i A_{i+1}$  representing  $C$  are linearly independent, a contradiction.

1606 We now show that both rows  $u_i$  and  $v_i$  in column  $e_i$  of  $T_i A_{i+1}$  are nonzero.  
 1607 Since  $e_i$  is not a loop of  $M$ , at least one entry of column  $e$  is nonzero; without  
 1608 loss of generality assume its entry in row  $u_i$  is not zero. For a contradiction,  
 1609 suppose its entry in row  $v_i$  is zero. Let  $Q$  be a subgraph of  $\Omega_i - v_i$  consisting of  
 1610 an unbalanced cycle and a path (possibly trivial) connecting this cycle to  $u_i$ .  
 1611 In  $F(\Omega_{i+1})$ ,  $E(Q) \cup e_i$  is independent, but the columns of  $T_i A_{i+1}$  representing  
 1612  $E(Q) \cup e_i$  are linearly dependent, a contradiction. Thus column  $e_i$  of  $T_i A_{i+1}$   
 1613 is nonzero in precisely its rows  $u_i, v_i$  corresponding to the endpoints of edge  $e_i$   
 1614 in  $\Omega_i$ . Let  $T_{i+1} = T_i$  and let  $S_{i+1}$  be the diagonal matrix obtained from  $S_i$  by  
 1615 adding a column to scale column  $e_i$  of  $T_{i+1} A_{i+1}$  so that its entry in row  $u_i$  is 1.  
 1616 Extend the  $\mathbb{F}^\times$ -gain function  $\varphi_i$  by defining  $\varphi_{i+1}(e_i)$  to be  $-(T_{i+1} A_{i+1} S_{i+1})_{v_i e}$ .  
 1617 Now  $T_{i+1} A_{i+1} S_{i+1}$  is the canonical frame matrix  $A_F(G_{i+1}, \varphi_{i+1})$ . By induction,  
 1618 there is a nonsingular matrix  $T_n$ , a diagonal matrix  $S_n$ , and a gain function  
 1619  $\varphi_n$  such that  $T_n A_n S_n = A_F(G_n, \varphi_n)$ .

1620 If  $(G, \mathcal{B})$  has no joints we are done. So assume  $J$  is nonempty. We now claim  
 1621 that  $M \neq L(G, \mathcal{B})$ . For suppose contrarily that  $M = L(G, \mathcal{B})$ . Since  $T_n A_n S_n$   
 1622 is a canonical frame representation of  $M \setminus J$ , it must be the case that  $\Omega_n$  is  
 1623 tangled. By Theorem 3.3  $\Omega_n$  contains a link minor  $(H, \mathcal{S})$  that is either a biased  
 1624  $2C_3$  with no balanced 2-cycle or a biased  $K_4$  with no balanced triangle. Since  
 1625  $J$  is nonempty  $(G, \mathcal{B})$  has a link minor  $(H', \mathcal{S}')$  where  $(H', \mathcal{S}')$  is obtained by  
 1626 adding a joint  $e$  incident to a vertex of  $(H, \mathcal{S})$ . Since  $T_n A_n S_n$  is a representation  
 1627 over  $\mathbb{F}$  for  $L(G, \mathcal{B})$  that agrees with  $T_n A_n S_n$  on all elements aside from possibly  
 1628 those in  $J$ , and since  $(H', \mathcal{S}')$  is a link minor of  $(G, \mathcal{B})$ , by Lemma 2.5 there  
 1629 is a matrix  $B$  over  $\mathbb{F}$  representing  $L(H', \mathcal{S}')$  with the property that  $B \setminus e$  is a  
 1630 canonical frame matrix particular to  $(H, \mathcal{S})$ . But this is impossible by Lemma  
 1631 5.8. Thus  $M \neq L(G, \mathcal{B})$ .

1632 Finally, we show that each column  $e$  of  $T_n A$  for which  $e \in J$  has exactly  
 1633 one nonzero entry. Suppose  $e \in J$  has endpoint  $u$  and that column  $e$  of  $T_n A$   
 1634 is nonzero in row  $v \neq u$ . Let  $C$  be the edge set of an unbalanced cycle in  
 1635  $\Omega_n - v$  together with a path linking this cycle and  $u$ . Then  $C$  is a circuit  
 1636 of  $F(G, \mathcal{B})$  but its corresponding columns in  $T_n A$  are linearly independent, a  
 1637 contradiction. Thus there is a diagonal matrix  $S$  scaling the columns of  $T_n A$   
 1638 such that  $T_n A S$  is a canonical frame matrix particular to  $(G, \mathcal{B})$ .

1639 (2) We proceed as in case (1), considering the matrix  $T_i A_{i+1}$ . Matrix  $A_{i+1}$   
 1640 has rows indexed by  $V(G) \cup v_0$ , according to the indexing of the corresponding

1641 rows of  $T_i A_i S_i$  where  $v_0 \notin V(G)$  corresponds to the “gains row” of  $T_i A_i S_i$ , and  
 1642 columns indexed by  $E(G_{i+1})$ . We first show that column  $e_i$  of  $T_i A_{i+1}$  is zero  
 1643 in every row aside from  $u_i$ ,  $v_i$ , and  $v_0$ . Suppose for a contradiction that  $e_i$  is  
 1644 nonzero in a row  $x \notin \{u_i, v_i, v_0\}$  of  $T_i A_{i+1}$ . As in case (1), let  $U_i \subseteq E(\Omega_i - x)$   
 1645 be a set of edges inducing a subgraph of  $\Omega_i$  that is a spanning tree of  $\Omega_i - x$   
 1646 along with one additional edge whose fundamental cycle with respect to this  
 1647 tree is unbalanced. Contained in  $U_i \cup e_i$  is a biased subgraph  $C$  whose edge set  
 1648 is a circuit of  $L(\Omega_{i+1})$ . The subgraph  $C$  contains  $e_i$  and does not contain  $x$ .  
 1649 Since  $T_i A_i S_i = A_L(G_i, \psi_i)$ , the columns of  $T_i A_{i+1}$  representing  $E(C) - e_i$  are  
 1650 all zero in row  $x$ . Hence while  $C$  is a circuit of  $L(\Omega_{i+1})$  the columns of  $T_i A_{i+1}$   
 1651 representing  $C$  are linearly independent, a contradiction.

1652 Since  $\Omega_i$  is unbalanced and 2-connected and  $\Omega_{i+1}$  is obtained from  $\Omega_i$  by  
 1653 adding a single edge,  $r(L(\Omega_{i+1})) = r(L(\Omega_i))$ . Thus  $A_i$  and  $A_{i+1}$  have the same  
 1654 rank. Since row  $v_0$  of  $T_i A_i S_i$  is not in the span of the rows  $V(G)$  neither is  
 1655 row  $v_0$  of  $T_i A_{i+1}$  in the span of the rows in  $V(G)$ . But the sum of the rows in  
 1656  $V(G)$  of  $T_i A_i$  is zero, so likewise the sum of the rows in  $V(G)$  of  $T_i A_{i+1}$  must  
 1657 be zero: otherwise the rank of  $T_i A_{i+1}$  would be greater than that of  $T_i A_i$ , a  
 1658 contradiction.

1659 Suppose first that both entries of column  $e$  in rows  $u_i$  and  $v_i$  are zero.  
 1660 Element  $e_i$  is not a loop of  $M$ , so then its entry in row  $v_0$  is nonzero. Let  $C$  be  
 1661 an unbalanced cycle in  $\Omega_i - u_i$ . Then  $C \cup e$  is independent in  $L(\Omega_{i+1})$  but the  
 1662 columns of  $T_i A_{i+1}$  representing  $C \cup e$  are linearly dependent, a contradiction.  
 1663 Thus column  $e_i$  has entries  $a$  and  $-a$  in rows  $u_i$  and  $v_i$ , where  $a \neq 0$ . Take  
 1664  $T_{i+1} = T_i$  and let  $S_{i+1}$  be the diagonal matrix obtained by adding a column  
 1665 to  $S_i$  to scale column  $e_i$  of  $T_{i+1} A_{i+1}$  by  $a^{-1}$ . Extend the  $\mathbb{F}^+$ -gain function  $\psi_i$   
 1666 to  $E(G_{i+1})$  by defining  $\psi(e_i)$  to be the entry in row  $v_0$  of column  $e_i$ . Now  
 1667  $T_{i+1} A_{i+1} S_{i+1}$  is the canonical lift matrix  $A_L(G_{i+1}, \psi_{i+1})$ . By induction, there  
 1668 is a nonsingular matrix  $T_n$ , a diagonal matrix  $S_n$ , and a gain function  $\psi_n$  such  
 1669 that  $T_n A_n S_n = A_L(G_n, \psi_n)$ .

1670 If  $(G, \mathcal{B})$  has no joints we are done. So assume  $J$  is nonempty. Analogous  
 1671 to the situation in case (1), we now claim that  $M \neq F(G, \mathcal{B})$ . Suppose to the  
 1672 contrary that  $M = F(G, \mathcal{B})$ . Since  $T_n A_n S_n$  is a canonical lift representation  
 1673 of  $M \setminus J$ , it must be the case that  $\Omega_n$  is tangled. By Theorem 3.3,  $\Omega_n$  contains  
 1674 a link minor  $(H, \mathcal{S})$  that is either a biased  $2C_3$  with no balanced 2-cycle or  
 1675 a biased  $K_4$  with no balanced triangle. Since  $J$  is nonempty  $(G, \mathcal{B})$  has a  
 1676 link minor  $(H', \mathcal{S}')$  where  $(H', \mathcal{S}')$  is obtained by adding a joint  $e$  incident to  
 1677 a vertex of  $(H, \mathcal{S})$ . Since  $T_n A_n S_n$  is a representation over  $\mathbb{F}$  for  $F(G, \mathcal{B})$  that  
 1678 agrees with  $T_n A_n S_n$  on all elements aside from possibly those in  $J$ , and since  
 1679  $(H', \mathcal{S}')$  is a link minor of  $(G, \mathcal{B})$ , by Lemma 2.5 there is a matrix  $B$  over  $\mathbb{F}$   
 1680 representing  $F(H', \mathcal{S}')$  with the property that  $B \setminus e$  is a canonical lift matrix

1681 particular to  $(H, \mathcal{S})$ . This violates Lemma 5.8, so  $M \neq F(G, \mathcal{B})$ .

1682 Finally, we show that each column  $e$  of  $T_n A$  for which  $e \in J$  has a nonzero  
 1683 entry only in row  $v_0$ . Suppose for a contradiction that  $e \in J$  with column  $e$  of  
 1684  $T_n A$  nonzero in row  $v \neq v_0$ . Let  $C$  be the edge set of an unbalanced cycle in  
 1685  $\Omega_n - v$ . Then  $C \cup e$  is a circuit of  $L(G, \mathcal{B})$  but its corresponding columns in  
 1686  $T_n A$  are linearly independent, a contradiction. Thus  $T_n A S_n$  is a canonical lift  
 1687 matrix particular to  $(G, \mathcal{B})$ .

1688 This completes the proof that at least one of statements (i) or (ii) of the  
 1689 theorem hold. But  $A_0$  is not projectively equivalent to both a canonical frame  
 1690 matrix and a canonical lift matrix particular to  $(G_0, \mathcal{B}_0)$ . Thus neither is  $A$   
 1691 projectively equivalent to both a canonical frame matrix and a canonical lift  
 1692 matrix particular to  $(G, \mathcal{B})$ .  $\square$

1693 For the second statement of Theorem 5.5, we would like to show that for  
 1694 each 3-connected matroid  $M$  represented by an almost-balanced biased graph  
 1695  $(G, \mathcal{B})$ , given any matrix  $A$  representing  $M$ ,  $A$  is projectively equivalent to a  
 1696 canonical lift matrix particular to  $(\widehat{G}, \widehat{\mathcal{B}})$ . Unfortunately, this can fail in the  
 1697 case  $M$  has rank 2. Let  $M$  be a 3-connected rank-2 matroid represented by  
 1698 the biased graph  $(G, \emptyset)$  consisting of  $E(M) - 1$  links between a pair of vertices  
 1699 and a single joint. Let  $\mathbb{F}$  be a field and let  $A$  be a matrix over  $\mathbb{F}$  representing  
 1700  $M$ . Then  $A$  is projectively equivalent to the matrix

$$1701 \quad A' = \begin{pmatrix} 1 & 0 & 1 & 1 & \cdots & 1 \\ 0 & 1 & a_1 & a_2 & \cdots & a_{n-2} \end{pmatrix}$$

1702 where  $n = |E(M)|$  and for each  $i$ ,  $a_i \notin \{0, 1\}$ . The matrix  $A'$  is a canonical  
 1703 frame representation particular to a roll-up of  $(G, \emptyset)$ , and  $A'$  is a canonical  
 1704 lift matrix particular to  $(G, \emptyset)$  (with its second row as the “gains row”). It is  
 1705 straightforward to see that by elementary row operations and column scaling  
 1706 we may obtain from  $A'$  a canonical frame matrix particular to  $(G, \emptyset)$ . How-  
 1707 ever,  $(\widehat{G}, \widehat{\emptyset})$  is the loopless contrabalanced biased graph obtained from  $(G, \emptyset)$   
 1708 by unrolling its joint, and there is no guarantee that  $A$  need be projectively  
 1709 equivalent to a canonical matrix representation particular to  $(\widehat{G}, \widehat{\emptyset})$ . Indeed,  
 1710 there is no guarantee that such a representation exists. For instance, if  $n = 6$   
 1711 and  $\mathbb{F}$  is  $\text{GF}(5)$ , then the above argument shows that  $A$  is projectively equiv-  
 1712 alent to a canonical frame matrix particular to a roll-up of  $(G, \emptyset)$ , and to a  
 1713 canonical lift matrix particular to  $(G, \emptyset)$ , but a canonical lift representation  
 1714 particular to  $(\widehat{G}, \widehat{\emptyset})$  requires six distinct elements of  $\mathbb{F}^+$ , so no such matrix  
 1715 exists. Fortunately, rank-2 is the only case in which this can occur:

1716 **Theorem 5.9.** *Let  $M$  be a matroid of rank greater than two represented by a*  
 1717 *2-connected almost-balanced biased graph  $(G, \mathcal{B})$  having no 2-separation with*  
 1718 *one side balanced. Let  $\mathbb{F}$  be a field and let  $A$  be an  $\mathbb{F}$ -matrix representing  $M$ .*  
 1719 *Then*

1720 (i)  *$A$  is projectively equivalent to a canonical lift matrix particular to  $(\widehat{G}, \widehat{\mathcal{B}})$ ,*  
 1721 *and*

1722 (ii)  *$A$  is projectively equivalent to a canonical frame matrix particular to each*  
 1723 *roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$ , unless  $\mathbb{F}$  is  $\text{GF}(2)$ .*

1724 *Furthermore,  $A$  is projectively equivalent to a canonical frame matrix partic-*  
 1725 *ular to  $(\widehat{G}, \widehat{\mathcal{B}})$  if and only if whenever  $\varphi$  is an  $\mathbb{F}^+$ -gain function for which  $A$*   
 1726 *and  $A_L(\widehat{G}, \varphi)$  are projectively equivalent,  $\varphi$  is switching equivalent to a gain*  
 1727 *function assigning 0 to  $e$  if and only if  $e$  is a link not incident to  $u$  or  $e$  is a*  
 1728 *loop of  $M$ .*

1729 We will need the following straightforward fact.

1730 **Lemma 5.10.** *Let  $C$  be the balanced triangle of  $D_{1,0}$  and let  $Y$  be a  $K_{1,3}$ -*  
 1731 *subgraph of  $D_{1,0}$  meeting  $C$  in exactly two edges. Then the simplification of*  
 1732  *$\nabla_Y D_{1,0}$  is isomorphic to  $B'_1$ .*

1733 *Proof of Theorem 5.9.* Suppose first that  $(\widehat{G}, \widehat{\mathcal{B}})$  does not contain a contrabal-  
 1734 *anced theta subgraph. The only biased graph representing  $U_{2,4}$  without a con-*  
 1735 *trabanced theta is shown at right in Figure 7. Moreover,  $F_7$  is neither frame*  
 1736 *nor lifted-graphic, and the only biased graphs representing  $F_7^*$ ,  $M^*(K_5)$ , and*  
 1737  *$M^*(K_{3,3})$  are all properly unbalanced [15]. Since none of these biased graphs*  
 1738 *can occur as a minor of  $(\widehat{G}, \widehat{\mathcal{B}})$  and  $(\widehat{G}, \widehat{\mathcal{B}})$  represents  $M$ ,  $M$  contains none of*  
 1739  *$U_{2,4}$ ,  $F_7$ ,  $F_7^*$ ,  $M^*(K_5)$ , nor  $M^*(K_{3,3})$  as a minor. Thus  $M$  is graphic. The first*  
 1740 *two statements follow. The third statement follows from Corollary 5.3.*

1741 So assume now that  $(\widehat{G}, \widehat{\mathcal{B}})$  contains a contrabanced theta subgraph. Let  
 1742  *$v$  be a balancing vertex of  $(\widehat{G}, \widehat{\mathcal{B}})$  and let  $J$  be the set of joints of  $(\widehat{G}, \widehat{\mathcal{B}})$ .*  
 1743 *Since the set of joints not incident to  $v$  form an unbalancing class of  $\Sigma(v)$*   
 1744 *in  $(G, \mathcal{B})$ , and all of the unbalancing classes of  $\Sigma(v)$  are unrolled in  $(\widehat{G}, \widehat{\mathcal{B}})$ ,*  
 1745 *every joint of  $(\widehat{G}, \widehat{\mathcal{B}})$  is incident to  $v$ . We claim that  $(\widehat{G}, \widehat{\mathcal{B}}) \setminus J$  does not have*  
 1746 *another balancing vertex  $u \neq v$ . For suppose to the contrary that  $(\widehat{G}, \widehat{\mathcal{B}}) \setminus J$*   
 1747 *has a balancing vertex  $u \neq v$ . Then  $(\widehat{G}, \widehat{\mathcal{B}}) \setminus J$  has the structure described in*  
 1748 *Proposition 2.12: there are graphs  $G_1, \dots, G_m$  such that  $\widehat{G} = G_1 \cup \dots \cup G_m$*   
 1749 *and  $G_j \cap G_k = \{u, v\}$  for each pair  $j \neq k$  and a cycle is in  $\widehat{\mathcal{B}}$  if and only if it is*  
 1750 *contained in a single graph  $G_j$ . Since  $(\widehat{G}, \widehat{\mathcal{B}})$  contains a contrabanced theta,*



1751  $m \geq 3$ . If there is a subgraph  $G_i$  with  $E(G_i) \geq 2$ , then  $(E(G_i), E(G) \setminus E(G_i))$  is  
 1752 a 2-separation of  $(G, \mathcal{B})$  with one side balanced, contrary to assumption. Thus  
 1753 for each  $i$ ,  $|E(G_i)| = 1$ . Thus  $G = mK_2$  and  $\mathcal{B}$  is empty. But  $M(mK_2, \emptyset)$  is  
 1754 isomorphic to the  $m$ -point line  $U_{2,m}$ , so  $r(M) = 2$ , contrary to assumption.

1755 Thus  $v$  is the unique balancing vertex of  $(\widehat{G}, \widehat{\mathcal{B}}) \setminus J$ . Since  $G$  is 2-connected,  
 1756 so is  $\widehat{G}$ . Thus by Proposition 3.5  $(\widehat{G}, \widehat{\mathcal{B}})$  contains a biased subgraph  $\Omega_0$  that  
 1757 is a subdivision of  $D_{1,0}$ ,  $B'_0$ ,  $B'_1$ , or  $B'_2$ . Since both  $\Omega_0$  and  $\widehat{G}$  are 2-connected,  
 1758 there is a sequence of 2-connected biased subgraphs  $\Omega_0 \subset \dots \subset \Omega_n$  where  
 1759  $\Omega_n = (\widehat{G}, \widehat{\mathcal{B}}) \setminus J$ , and for each  $i \in \{0, \dots, n-1\}$  there is a path  $P_i$  in  $G$   
 1760 internally disjoint from  $\Omega_i$  so that  $\Omega_{i+1} = \Omega_i \cup P_i$ . For each  $i \in \{0, \dots, n\}$ , let  
 1761  $(G_i, \mathcal{B}_i)$  be the biased subgraph of  $(\widehat{G}, \widehat{\mathcal{B}})$  induced by  $E(\Omega_i)$  with  $V(G_i) = V(\widehat{G})$   
 1762 and let  $A_i$  be the submatrix of  $A$  consisting of all rows of  $A$  and precisely  
 1763 those columns representing  $E(\Omega_i)$ . Thus for each  $i$ ,  $A_i$  represents  $M(\Omega_i)$ . By  
 1764 Lemmas 5.7 and 4.19, Proposition 2.9, and Lemma 5.10,  $A_0$  is projectively  
 1765 equivalent to a canonical lift matrix particular to  $\Omega_0$ . Inductively assume that  
 1766 there are sequences  $T_0, \dots, T_i$  and  $S_0, \dots, S_i$  of  $\mathbb{F}$ -matrices such that for each  
 1767  $k \in \{0, \dots, i\}$  each matrix  $T_k A_k S_k$  is a canonical lift matrix particular to  $\Omega_k$ .  
 1768 We may assume that  $A$  is of full rank, and so that for each  $k \in \{0, \dots, i\}$  the  
 1769 rows of  $T_k A_k S_k$  are indexed by  $v_0 \cup (V(G_k) - v)$  (as described in Section 2.6).  
 1770 Since for each  $k$ ,  $V(G_k) = V(\widehat{G})$ , for each  $k$  the rows of  $T_k A_k S_k$  are indexed  
 1771 by  $v_0 \cup (V(\widehat{G}) - v)$ .

1772 Consider  $\Omega_{i+1} = \Omega_i \cup P_i$  and the matrix  $T_i A_{i+1}$  representing  $M(\Omega_{i+1})$ . Let  
 1773 us assume  $P_i$  consists of a single edge  $e_i$  whose endpoints are  $x_i, y_i \in V(\Omega_i)$ .  
 1774 Let  $x \in V(G_i) - \{x_i, y_i\}$ . Since  $\Omega_i$  is 2-connected, there is a spanning tree  $T$  of  
 1775  $\Omega_i - x$ . If  $x$  is not  $v$ , then there is an edge  $f$  such that the fundamental cycle in  
 1776  $T \cup f$  is unbalanced in  $\Omega_i$ : set  $W = E(T) \cup f$ ; otherwise set  $W = E(T)$ . The  
 1777 subgraph  $W \cup e_i \subseteq \Omega_{i+1}$  contains a subgraph  $C$  that is either a balanced cycle,  
 1778 a pair of unbalanced cycles sharing just vertex  $v$ , or a contrabalanced theta.  
 1779 Since  $E(C)$  is a circuit of  $M(\Omega_{i+1})$ , this implies that in column  $e_i$  of  $T_i A_{i+1}$ , the  
 1780 entry in row  $x$  is 0. As long as neither endpoint of  $e_i$  is  $v$ , this also implies that  
 1781 the entries in rows  $x_i$  and  $y_i$  are nonzero. If  $e_i$  has endpoints  $v, y_i$ , for some  
 1782 vertex  $y_i \neq v$ , then the form of  $C$  in  $\Omega_{i+1}$  implies that the entry in matrix  
 1783  $T_i A_{i+1}$  in column  $e_i$ , row  $y_i$  must be nonzero. Thus  $T_i A_{i+1} S_{i+1}$ , with rows  
 1784 indexed by  $v_0 \cup (V(\widehat{G}) - v)$ , is a canonical lift matrix for some appropriate  
 1785 column scaling matrix  $S_{i+1}$ . Hence by induction, there are matrices  $T_n$  and  $S_n$   
 1786 such that  $T_n A_n S_n$  is a canonical lift matrix particular to  $\Omega_n = (\widehat{G}, \widehat{\mathcal{B}}) \setminus J$ .

1787 Finally, consider the set of joints  $J$ . Let  $e$  be a joint. By assumption  $e$  is  
 1788 incident to  $v$  and every other joint is in parallel with  $e$ . Since  $v$  is the unique  
 1789 balancing vertex of  $(\widehat{G}, \widehat{\mathcal{B}})$ , for every vertex  $x \neq v$ , there is an unbalanced cycle

1790  $C_x$  in  $G - x$  of length  $> 1$ . Since  $C_x \cup e$  is a circuit of  $M$ , row  $x$  of column  $e$   
 1791 of  $T_n A$  is zero. Thus every row of column  $e$  aside from row  $v_0$  is zero. Since  
 1792  $e$  is not a loop of  $M$ , the entry in row  $v_0$  of column  $e$  must be nonzero. Since  
 1793 all joints of  $(\widehat{G}, \widehat{\mathcal{B}})$  are in parallel with  $e$ , every column of  $A$  representing a  
 1794 joint is zero in all rows but  $v_0$ . Thus there is a diagonal matrix  $S$  scaling the  
 1795 columns of  $T_n A$  so that  $T_n AS$  is a canonical lift matrix particular to  $(G, \mathcal{B})$ .  
 1796 This completes the proof of statement (i).

1797 (ii) Let  $(H, \mathcal{S})$  be a roll-up of  $(\widehat{G}, \widehat{\mathcal{B}})$ , and let  $U \subseteq \Sigma(v)$  be the unique  
 1798 unbalancing class of edges in  $\Sigma(v)$  that are joints in  $(H, \mathcal{S})$ . By statement  
 1799 (i) there is a  $\mathbb{F}^+$ -gain function  $\varphi$  on  $\widehat{G}$  realizing  $\widehat{\mathcal{B}}$  for which  $A = A_L(\widehat{G}, \varphi)$ .  
 1800 Let  $T$  be a spanning tree of  $\widehat{G}$  containing exactly one edge in  $U$ . Then the  
 1801  $T$ -normalized gain function  $\varphi'$  obtained by switching on  $\varphi$  satisfies  $\varphi'(e) = 0$   
 1802 for all edges  $e$  not incident to  $v$  and  $\varphi'(e) = 0$  for each edge in  $U$ . By Corollary  
 1803 5.3, the derived  $\mathbb{F}^\times$ -gain function  $\varphi'^\times$  realizes  $(H, \mathcal{S})$ . By Lemma 5.2,  $A$  and  
 1804  $A_F(H, \varphi'^\times)$  are projectively equivalent.

1805 The final statement follows immediately from Corollary 5.3.  $\square$

### 1806 5.3 Projective equivalence classes are in 1-1 correspon- 1807 dence with switching classes

1808 Finally, we can prove Theorem 3.

1809 *Proof of Theorem 3.* By Proposition 4.2, every switching class of gain func-  
 1810 tions is contained in a projective equivalence class of matrix representations.  
 1811 Thus we just need show that if  $A$  and  $B$  are projectively equivalent repre-  
 1812 sentations of  $M$ , then  $A$  and  $B$  are each projectively equivalent to canonical  
 1813 representations whose gain functions are contained in the same switching class.  
 1814 Because  $M$  is 3-connected,  $(G, \mathcal{B})$  is 2-connected and has no 2-separation with  
 1815 one side balanced.

1816 Assume first that  $(G, \mathcal{B})$  is properly unbalanced and not tangled. Then ei-  
 1817 ther  $M = F(G, \mathcal{B})$  or  $M = L(G, \mathcal{B})$ , but not both. Assume that  $M = F(G, \mathcal{B})$ .  
 1818 Let  $A$  and  $B$  be projectively equivalent  $\mathbb{F}$ -matrices representing  $M$ . By The-  
 1819 orem 5.5(1), each of  $A$  and  $B$  are projectively equivalent to a canonical frame  
 1820 matrix particular to  $(G, \mathcal{B})$ . Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^\times$ -gain functions such that  
 1821  $A$  is projectively equivalent to  $A_F(G, \varphi)$  and  $B$  is projectively equivalent to  
 1822  $A_F(G, \psi)$ . By Theorem 5.1(1),  $\varphi$  and  $\psi$  are switching equivalent. Similarly, if  
 1823  $M = L(G, \mathcal{B})$  and  $A$  and  $B$  are projectively equivalent  $\mathbb{F}$ -matrices representing  
 1824  $M$ , then by Theorem 5.5(1), each of  $A$  and  $B$  are projectively equivalent to a  
 1825 canonical lift matrix particular to  $(G, \mathcal{B})$ . Let  $\varphi$  and  $\psi$  be  $\mathbb{F}^+$ -gain functions

1826 such that  $A$  is projectively equivalent to  $A_L(G, \varphi)$  and  $B$  is projectively equiv-  
 1827 alent to  $A_L(G, \psi)$ . By Theorem 5.1(2),  $\varphi$  and  $\psi$  are switching-and-scaling  
 1828 equivalent.

1829 Now assume that  $(G, \mathcal{B})$  is properly unbalanced but tangled. Then  $L(G, \mathcal{B})$   
 1830 and  $F(G, \mathcal{B})$  coincide:  $M = L(G, \mathcal{B}) = F(G, \mathcal{B})$ . Let  $A$  and  $B$  be projectively  
 1831 equivalent  $\mathbb{F}$ -matrices representing  $M$ . By Theorem 5.5(1), each of  $A$  and  $B$  is  
 1832 projectively equivalent to a canonical lift matrix particular to  $(G, \mathcal{B})$ , or to a  
 1833 canonical frame matrix particular to  $(G, \mathcal{B})$ , but not both. By Theorem 5.1(3),  
 1834 either  $A$  and  $B$  are both projectively equivalent to canonical lift matrices or  
 1835 both are projectively equivalent to canonical frame matrices. In either case,  
 1836 by statement (1) or (2) of Theorem 5.1, the gain functions from which these  
 1837 canonical representations arise belong to the same switching class.

1838 Finally, assume that  $(G, \mathcal{B})$  is almost-balanced, and let  $A$  and  $B$  be projec-  
 1839 tively equivalent  $\mathbb{F}$ -matrices representing  $M$ . By Proposition 2.12,  $(G, \mathcal{B})$  has  
 1840 a unique balancing vertex. By Theorem 5.5(2),  $A$  and  $B$  are each projectively  
 1841 equivalent to a canonical lift matrix particular to  $(\widehat{G}, \widehat{\mathcal{B}})$ , say, given by  $\mathbb{F}^+$ -gain  
 1842 functions  $\varphi$  and  $\psi$  respectively. By Theorem 5.4(1),  $\varphi$  and  $\psi$  belong to the  
 1843 same switching class.  $\square$

## 1844 References

- 1845 [1] N. Bowler, D. Funk, and D. Slilaty, *Describing Quasi-Graphic Matroids*,  
 1846 ArXiv e-prints (2018).
- 1847 [2] Tom Brylawski, *Modular constructions for combinatorial geometries*,  
 1848 Trans. Amer. Math. Soc. **203** (1975), 1–44.
- 1849 [3] R. Chen and I. Pivotto, *Biased graphs with no two vertex-disjoint unbal-  
 1850 anced cycles*, ArXiv e-prints (2014).
- 1851 [4] Henry H. Crapo, *Single-element extensions of matroids*, J. Res. Nat. Bur.  
 1852 Standards Sect. B **69B** (1965), 55–65. MR 0190045
- 1853 [5] Matt DeVos and Daryl Funk, *Almost balanced biased graph represen-  
 1854 tations of frame matroids*, Advances in Applied Mathematics **96** (2018),  
 1855 139–175.
- 1856 [6] Jim Geelen, *Some open problems on excluding a uniform matroid*, Adv.  
 1857 in Appl. Math. **41** (2008), no. 4, 628–637. MR 2459453 (2009k:05050)

- 1858 [7] Jim Geelen, Bert Gerards, and Geoff Whittle, *Quasi-graphic matroids*,  
1859 arXiv:1512.03005, 14pp.
- 1860 [8] ———, *The highly connected matroids in minor-closed classes*, *Ann.*  
1861 *Comb.* **19** (2015), no. 1, 107–123.
- 1862 [9] J. Kahn and J. P. S. Kung, *Varieties of combinatorial geometries*, *Trans.*  
1863 *Amer. Math. Soc.* **271** (1982), no. 2, 485–499. MR 654846 (84j:05043)
- 1864 [10] James Oxley, *Matroid theory*, second ed., Oxford Graduate Texts in Math-  
1865 ematics, vol. 21, Oxford University Press, Oxford, 2011. MR 2849819  
1866 (2012k:05002)
- 1867 [11] Daniel Slilaty, *Projective-planar signed graphs and tangled signed graphs*,  
1868 *J. Combin. Theory Ser. B* **97** (2007), no. 5, 693–717. MR 2344133  
1869 (2008j:05161)
- 1870 [12] Geoff Whittle, *Stabilizers of classes of representable matroids*, *J. Combin.*  
1871 *Theory Ser. B* **77** (1999), no. 1, 39–72. MR 1710531 (2000f:05027)
- 1872 [13] Thomas Zaslavsky, *Vertices of localized imbalance in a biased graph*, *Proc.*  
1873 *Amer. Math. Soc.* **101** (1987), no. 1, 199–204. MR 897095 (88f:05103)
- 1874 [14] ———, *Biased graphs. I. Bias, balance, and gains*, *J. Combin. Theory*  
1875 *Ser. B* **47** (1989), no. 1, 32–52. MR 1007712 (90k:05138)
- 1876 [15] Thomas Zaslavsky, *Biased graphs whose matroids are special binary ma-*  
1877 *troids*, *Graphs Combin.* **6** (1990), no. 1, 77–93. MR 1058551 (91f:05097)
- 1878 [16] Thomas Zaslavsky, *Biased graphs. II. The three matroids*, *J. Combin.*  
1879 *Theory Ser. B* **51** (1991), no. 1, 46–72. MR 1088626 (91m:05056)
- 1880 [17] Thomas Zaslavsky, *Frame matroids and biased graphs*, *European J. Com-*  
1881 *bin.* **15** (1994), no. 3, 303–307. MR 1273951 (95a:05021)
- 1882 [18] ———, *Biased graphs. IV. Geometrical realizations*, *J. Combin. Theory*  
1883 *Ser. B* **89** (2003), no. 2, 231–297. MR 2017726 (2005b:05057)